

Efficacy of Photobiomodulation Therapy in the Treatment of Pain and Inflammation: A Literature Review

As a library, NLM provides access to scientific literature. Inclusion in an NLM database does not imply endorsement of, or agreement with, the contents by NLM or the National Institutes of Health.

[Ana González-Muñoz](#),^{1,2} [María Cuevas-Cervera](#),¹ [José Javier Pérez-Montilla](#),¹ [Daniel Aguilar-Núñez](#),³ [Dina Hamed-Hamed](#),¹ [María Aguilar-García](#),¹ [Leo Pruiomboom](#),⁴ and [Santiago Navarro-Ledesma](#)^{1,4,*}

George A. Koumantakis, Academic Editor

Author information Article notes Copyright and License information [Disclaimer](#)

Abstract

The main objective of this literature review was to analyze the efficacy of (PBM) therapy application on subjects with chronic pain and inflammation, and furthermore, to evaluate

the methodological quality of the collected literature. The search was conducted using five databases: PubMed, ProQuest, Scopus, Web of Science, and PEDro. The keywords "low level laser therapy", "chronic pain", and "inflammation" provided the selection of RCTs that were published within the last 5 years, conducted in humans, and written in English. The PEDro Internal Validity Scale (IVS) checklist was used to evaluate the risk of bias in the included studies. A total of 11 articles were selected, all of them RCTs. Of the articles, five showed that PBM positively influences chronic pain, while another showed the same but only in the short term. In two other articles, the patient's inflammation improved markedly. In one article there was no improvement in chronic pain and in another, there was no improvement in inflammation. Four articles demonstrated that PBM is beneficial in acute pain. Furthermore, six studies were given an "excellent" score and the remaining five a "good" score based on the IVS. Photobiomodulation has beneficial effects on chronic pain and inflammation, although more research needs to be completed in this line for this to be clarified as the existence of RCTs on this subject is limited.

Keywords: low-level laser therapy, photobiomodulation, chronic pain, inflammation, metabolism

1. Introduction

Photobiomodulation therapy (PBM) commonly uses

wavelengths of light with an energy density ranging from 1 to 150 J/cm² and from 600 to 1070 nm. The effective tissue penetration is maximal in this range with hemoglobin and melanin, as the principal tissue chromophores, having high absorption bands at wavelengths shorter than 600 nm. The treatment of superficial tissue uses wavelengths in the range of 600 to 700 nm, while the treatment of deeper tissue uses wavelengths in the range of 780 to 950 nm [1,2]. Currently, whole-body PBM has shown a systemic response in addition to the local response, with improvements in quality of life, pain, sleep disorders, tiredness, muscle spasm, morning stiffness, psychological factors, elastic properties of tissue, circadian rhythms, tender points, and in fibromyalgia sufferers [3,4,5,6,7].

Furthermore, PBM therapy has also been shown to improve cerebral blood flow, neuronal bioenergetic functions, neuroinflammation, oxidative stress, neural apoptosis, neurogenesis, and neurotrophic factors, and additionally has effects on intrinsic brain networks [1]. In addition, PBM is also thought to affect the secretion of certain hormones, such as serotonin and endorphins, leading to a reduction in pain signaling [8].

Chronic pain is a problem that has a very notorious impact on society and people's lives and is one of the most common health problems among older adults (>65 years). It is

estimated that 13–50% of adults in the United Kingdom suffer from chronic pain although it is difficult to obtain accurate data since the estimates for the prevalence in the population vary greatly by place, time, and population. It has been estimated to be 8% per year in the UK. On the other hand, the US sets its estimated costs attributable to chronic pain, including disability, loss of work, and treatments, at about USD 600 million annually, with an incidence of 28.4% in the adult population. Treatment in these older adults is complex since it must have a multifactorial focus that includes pharmacological interventions, physical rehabilitation, and procedures to eradicate the cycle of pain [9,10,11,12]. By definition, chronic pain is pain that persists for at least 3 months. It is a factor in premature death and accelerated cognitive deterioration. In turn, this deterioration and dementia make treatment decisions difficult since the patient's ability to perceive pain and report it is impaired [11].

Inflammation is present in many chronic pain conditions and its reduction is usually related to a decrease in pain.

Oxidative stress is known to contribute to this inflammation [12]. Prolonged inflammation can lead to chronic hypersensitivity and perpetuate chronic pain. Several factors have been identified that influence inflammation induced by oxidative stress, chemokines, transcription factors, microRNAs, and inflammatory cytokines [12]. It has also been proven that, if widespread inflammation in neurological

tissue, which has been caused by a failure of the immune system to promote homeostasis, lasts beyond a certain time, it can result in the onset of chronic pain. When this failure continues, low-grade inflammation (LGI) and existing damage develop, and its perpetuation indicates the development of chronic musculoskeletal pain, in addition to peripheral hypersensitization, which evolves into central hypersensitization. Therefore, LGI is thought of as a risk factor for chronic diseases.

Photobiomodulation has been used in pain and inflammation treatment; nevertheless, a synthesis of its effects and the quality analysis of the studies have never been studied. The goal of this study was to analyze the scientific evidence that currently exists on PBM's effectiveness when used in the treatment of chronic musculoskeletal pain and inflammation. Secondly, to evaluate the methodological quality of the collected literature.

2. Methodology

2.1. Study Design

The present study is a literature review on randomized clinical trials (RCTs). In order to achieve a high methodological quality, a methodology characteristic of a systematic review, including the methodological quality of the selected RCTs, was carried out in accordance with the

PRISMA criteria for the development and preparation of meta-analysis and systematic reviews [13].

2.2. Sources of Information and Search Strategies

The literature search was carried out between March, April, and May 2022 using the following five electronic databases: PubMed, ProQuest, Web of Science, Scopus, and PEDro (see [Table 1](#)).

Table 1

Search strategy in the different databases used.

PubMed	<ul style="list-style-type: none">• "Low-Level Light Therapy" (MeSH) AND "chronic pain" (MeSH)• "Low-level Laser Therapy" (MeSH) OR "photobiomodulation" AND "physiotherapy"• "Low-level Light Therapy" (MeSH) AND "inflammation" (MeSH)
ProQuest	<ul style="list-style-type: none">• "Low-Level Light Therapy" AND "chronic pain"• "Low-level Light Therapy" OR "photobiomodulation" AND "physiotherapy"• "Low-level Light Therapy" AND "metabolism"• "Low-level Light Therapy" AND "inflammation"

Scopus	<ul style="list-style-type: none"> • "Low-level light therapy" AND "chronic pain" • "Low-level light therapy" AND "inflammation"
Web of Science	<ul style="list-style-type: none"> • "Low-level light therapy" AND "chronic pain" • "Low-level light therapy" AND "inflammation"
PEDro	<ul style="list-style-type: none"> • "Low-level light therapy" AND "chronic pain" • "Low-level light therapy" AND "inflammation"

The PICO strategy components were followed to develop the research questions and were based on the following elements:

- Population: patients older than 18 years of age and with inflammation or pain lasting for more than 3 months.
- Intervention: reduction in the symptomology of pain or inflammation after a PBM intervention.
- Comparison: the use or not of PBM.
- Results: effectiveness of the PBM application strategies on chronic pain and inflammation.

2.2.1. PubMed

The search for articles in PubMed used Medical Subject Headings (MeSH), as well as key terms not included in

MeSH such as “photobiomodulation”.

The keywords were “Low-level laser therapy”, “metabolism”, “inflammation”, “physiotherapy”, “chronic pain” and “photobiomodulation”.

The Boolean operators were: AND/OR.

The following parameters were used to limit the search strategy of this study:

- Text accessibility: open-access or accessible to the University of Granada, Spain.
- Type of articles: RCTs and observational studies.
- Date of publication: 2017–2022.
- Species: human.
- Language: English.

2.2.2. ProQuest

The keywords “low-level laser therapy”, “metabolism”, “photobiomodulation”, “inflammation”, “physiotherapy” and “chronic pain” were used for the search.

The Boolean operators were: AND/OR.

The following parameters were used to limit the search

strategy of this study:

- Text accessibility: open-access or accessible to the University of Granada, Spain.
- Type of articles: RCTs and observational studies.
- Date of publication: 2017–2022.
- Language: English.

2.2.3. Scopus

The keywords used for the search were: "Low-level laser therapy", "chronic pain", "photobiomodulation" "physiotherapy, "metabolism" and "inflammation".

The Boolean operator used was: AND.

The following parameters were used to limit the search strategy of this study:

- Text accessibility: open-access or accessible to the University of Granada, Spain.
- Type of articles: articles.
- Subject: medicine and health professions.
- Date of publication: 2017–2022.

2.2.4. Web of Science

The keywords for the search were: "Low-level laser therapy", "chronic pain", "photobiomodulation", "metabolism", "physiotherapy" and "inflammation".

The Boolean operator was: AND.

The following parameters were used to limit the search strategy of this study:

- Text accessibility: open-access or accessible to the University of Granada, Spain.
- Type of articles: articles (RCTs).
- Subject: medicine and health professions.
- Date of publication: 2017–2022.

2.2.5. PEDro

The keywords for the search were: "Low-level laser therapy", "chronic pain", "physiotherapy", "photobiomodulation", "metabolism" and "inflammation".

The Boolean operator was: AND.

The following parameters were used to limit the search strategy:

- Text accessibility: open-access or accessible to the University of Granada, Spain.
- Subject: chronic pain.
- Method: RCTs.
- Date of publication: 2017–2022.

2.3. Study Selection

2.3.1. Criteria for Inclusion

- Publication between January 2017 and May 2022.
- International publication in English and peer-reviewed.
- Designed as RCTs.
- Access to the complete text available.
- Conducted in humans over the age of 18.
- Suffer from acute pain or chronic pain.
- Suffer from some type of pathology that causes chronic pain.
- Suffer from edema/inflammation.

2.3.2. Exclusion Criteria

- Not meeting the inclusion criteria.
- Repeated in the databases.
- Does not assess chronic pain, inflammation or pathology or symptom prevention.
- Lack of PBMT implementation in people with pain or inflammation.
- No results were shown or no interpretation of the data.

2.4. Article Selection

This literature review aimed at focusing on photobiomodulation therapy in general. For this reason, the scope of the search included all areas of low-intensity laser research and the impact it has on various facets within the health fields of medicine and physiotherapy. These studies encompassed the effect of a low-intensity laser on diseases such as fibromyalgia to articles that revealed the part these lasers play in impairments in our temporomandibular joint or teeth. The first searches resulted in 59 eligible studies, which either proved or linked PBM and the benefits it can bring to health regarding the most important areas we wanted to cover in this study.

In order to increase the search quality, more focused searches were carried out on the influence PBM has on

chronic pain, at all levels, without neglecting the impact it has on the diseases and injuries in the subjects studied.

Although the search terms were further limited and reflected scientific evidence for PBM in subjects over 18 years of age, studies that did not fall within the scope of our study continued to appear.

Photobiomodulation therapy has been classified as a safe, non-invasive treatment modality. Several possible mechanisms have been attributed to PBM such as an increased thermal threshold, improved blood circulation, increased oxygen consumption by accelerating the redox reaction rate, and increased production of ATP and proinflammatory cytokines [14].

Therefore, the scope of this literature review was reduced to studies that researched any link between PBM and the benefits it can bring to our health, with the primary focus on its influence on pain and inflammation.

2.5. Methodological Quality Assessment of the RCTs

The assessment of the methodological quality of the selected RCTs followed the PEDro scale, which was adapted and translated to the Spanish language and allowed the items to be both validated and evaluated.

Each item, excluding the first, raises the overall score by 1 point. Eleven items can be included, resulting in a global score of 0 to 10 points [15].

Subsequently, the Internal Validity Score (IVS) was used to evaluate the internal validity of the RCTs. This scale contains the items that are indicated to be the most representative in ascertaining the internal validity of an article. The selected items are 2, 3, 5, 6, 7, 8, and 9, so the internal validity will have a score of up to 7 points [15].

2.6. RCTs Qualitative Synthesis

The score given by the scale determined the quality of the study. Hence, a score of 9–10 points is accepted as having excellent methodological quality, a score of 6–8 points has a good methodological quality, those in the 4–5 point range have a regular quality, and finally, those trials that obtain less than 4 points have a poor methodological quality. Studies that fall within this last range were discarded from this study [15].

The IVS classification of the studies is as follows: studies that receive 6–7 points are given a methodological quality that is high, those with 4–5 points a moderate methodological quality, and 0–3 points have a limited methodological quality [15].

2.7. Management of the Identified Literature

The bibliographical references that appear in this review have been managed using Mendeley software.

3. Results

This section provides the results found in the selected RCTs.

3.1. Study Selection

The searches resulted in 96,751 articles being found in the 5 databases as follows: PubMed, Scopus, PEDro, ProQuest, and Web of Science.

Articles that did not meet the inclusion criteria set out in the "inclusion criteria" section were discarded. The titles and summaries of the found articles were read and the information was summarized to highlight the most relevant aspects, both positive and negative, of this therapy.

The end result of the search and the methodological quality study of the preselected articles was 11 RCTs that met the admission criteria. The RCT selection process is shown in [Figure 1](#) by means of a flowchart in accordance with the guidelines and criteria established in the PRISMA 2009 declaration [13].

[Figure 1](#)

M. et al., 2018 [20]	1	1	1	1	1	1	1	1	1	1	1	10/10 Excellent
De Souza R. et al., 2018 [21]	1	1	0	0	0	0	1	1	1	1	1	6/10 Good
Shaiane S. et al., 2021 [22]	1	1	1	1	1	1	1	0	1	1	1	9/10 Excellent
Chong R. et al., 2020 [23]	1	1	1	1	1	1	1	1	1	1	1	10/10 Excellent
Taradaj J. et al., 2019 [24]	1	1	1	1	1	0	1	1	1	1	1	9/10 Excellent
Fang-Yin L. et al., 2020 [25]	1	1	1	1	1	0	1	1	0	1	1	8/10 Good

1. The selection criteria were specified. 2. Subjects were randomly assigned to groups. 3. Allocation was concealed. 4. Groups were similar at baseline regarding the most important prognostic indicators. 5. All subjects were blinded. 6. All therapists who administered the therapy were blinded. 7. All assessors who measured at least one key outcome were blinded. 8. Measures of at least one of the key outcomes were obtained from more than 85% of the subjects initially assigned to the groups. 9. Results were

presented for all subjects who received treatment or were allocated to the control group, or where this could not be, data for at least one key outcome was analyzed by “intention to treat”. 10. The results of statistical comparisons between groups were reported for at least one key outcome. 11. The study provides point and variability measures for at least one key outcome. * Non-summation criterion for the PEDro scale [15].

Table 3

Internal validity of the selected RCTs.

Author, Year	2	3	5	6	7	8	9	IVS
Eduardo C. et al., 2020 [16]	1	1	1	1	0	1	1	6/7 High
Luciana GL. et al., 2018 [17]	1	1	1	1	1	1	1	7/7 High
Fernanda R. et al., 2018 [18]	1	0	1	1	0	0	0	3/7 Limited
LeylaK. et al., 2020 [19]	1	1	1	0	0	0	0	3/7 Limited
Mariana M. et al., 2018 [20]	1	1	1	1	1	1	1	7/7 High
De Souza R. et al., 2018 [21]	1	0	0	0	1	1	1	4/7 Limited
Shaiane S. et al., 2021 [22]	1	1	1	1	1	0	1	6/7 High

Chong R. et al., 2020 [23]	1	1	1	1	1	1	1	7/7 High
Taradaj J. et al., 2019 [24]	1	1	1	0	1	1	1	6/7 High
Fang-Yin L. et al., 2020 [25]	1	1	1	0	1	1	0	5/7 High
Fernanda T. et al., 2018 [26]	1	0	0	0	1	1	1	4/7 Limited

IVS: Internal Validity Score [15].

Table 4

Development of the general characteristics for the selected studies.

Author, Year	Type of Study	Experimental Group	Control Group
Nunes EC. et al., 2020 [16]	Randomized Clinical Trial	<p>35 people with post endodontic surgery pain.</p> <p>Men: 16 Women: 19</p> <p>Age range: 20.8–37.2</p> <p>Chronic illness: No</p> <p>Post operative</p>	<p>35 people with post endodontic surgery pain.</p> <p>Men: 15 Women: 20</p> <p>Age range: 18.4–42.2</p> <p>Chronic illness: No</p> <p>Post</p>

		<p>pain: 1.3–2.9</p> <p>Treatment time: 39.2–54.2</p>	<p>operative pain: 1.4–3.2</p> <p>Treatment time: 49.9– 39.3</p>
<p>Gonçalves Langella L. et al., 2018 [17]</p>	<p>Randomized Clinical Trial</p>	<p>9 people in the post-surgical period of knee arthroplasty.</p>	<p>9 people in the post- surgical period of knee arthroplasty.</p>
<p>Fernanda R. et al., 2018 [18]</p>	<p>Randomized Clinical Trial</p>	<p>Group US + LLLT: 14 Caucasian women with knee osteoarthritis.</p> <p>Age range: 77–67</p> <p>Group US + ET + LLLT: 14 Caucasian women with knee</p>	<p>14 Caucasian women with knee osteoarthritis.</p> <p>Age range: 69–61.</p>

osteoarthritis.

Age range: 71–61

Leyla K. et al., 2020 [19]

Randomized Clinical Trial

20 people with nonspecific chronic low back pain.

Men: 7
Women: 13

Age range: 25–65
Average age: 47

20 people with nonspecific chronic low back pain.

Men: 5
Women: 15

Age range: 28–69

Average age: 51

<p>Mariana M. et al., 2017 [20]</p>	<p>Randomized Clinical Trial</p>	<p>PBMT group: Formed of 2 groups, each containing 20 people with FM.</p> <p>ET group: Formed of 2 groups, each containing 20 people with FM.</p> <p>PBMT + ET group: Formed of 2 groups, each containing 20 people with FM.</p> <p>Set 1: Investigates the immediate effect of a single PBMT/ET session on chronic pain condition Age: 32–38 BMI (Kg/m²): 21–31</p> <p>Set 2: Investigates the long-term</p>	<p>Divided into 2 groups, each containing 20 people with FM.</p> <p>Set 1: Investigates the immediate effect of a single PBMT/ET session on chronic pain condition Age: 32–38 BMI (Kg/m²): 21–31</p> <p>Set 2: Investigates the long-term effect (10 weeks) on the chronic pain condition and</p>

		<p>effect (10 weeks) on the chronic pain condition and other FM symptoms</p> <p>Age: 38–42 BMI(Kg/m²): 23–31</p>	<p>other FM symptoms</p> <p>Age: 38–42 BMI (Kg/m²): 23–31</p>
De Souza R. et al., 2018 [21]	Randomized Clinical Trial	<p>66 people with FM.</p> <p>Age Range: 57.05–35.23 Average age: 46.14</p> <p>(62 women and 4 men)</p> <p>Group A: 33 volunteers with FM and chronic pain.</p> <p>Group B: 33 volunteers with FM</p>	None.

and chronic pain.

Shaiane
S. et al.,
2021 [22]

Randomized
Clinical Trial

9 people with
chronic nonspecific
low back pain.

Women: 5
Men: 4

Age range: 20.51–
47.05
Average age: 33.78

9 people with
chronic
nonspecific
low back
pain.

Women: 6
Men: 3

Age range:
19.63–44.23
Average age:
31.44

27 people with

27 people
with

Chong R.
et al.,
2020 [23]

Randomized
Clinical Trial

orthodontic pain
and periodontal
inflammation.

orthodontic
pain and
periodontal
inflammation.

Taradaj J.
et al.,
2019 [24]

Randomized
Clinical Trial

TLAI group: 18
people with
nonspecific chronic
low back pain.

Men: 10

Women: 8

Age range: 29–58.

Average age: 44

LLLT group: 16
people with
nonspecific chronic
low back pain.

Men: 8

Women: 8

Age range: 29–53

Average age: 45.50

TLAI group:
17 people
with

nonspecific
chronic low
back pain.

Men: 9

Women: 8

Age range:
26–51.

Average age:
45

LLLT group:
17 people
with

nonspecific
chronic low
back pain.

Men: 9

Women: 8

Age range:
22–76

Average age:
52

Fang-Yin
L. et al.,
2020 [25]

Randomized
Clinical Trial

16 people with
knee osteoarthritis.

Age range: 77.42–
63.64.

17 people
with knee
osteoarthritis.

Age range:
76.64–62.82.

Fernanda T. et al., 2018 [26]	Randomized Clinical Trial	<p>Groups made up of 41 patients with temporomandibular pain and dysfunction (TMD).</p> <p>PBMT group: 14 patients. Women: 14 Age range: 30–61.4</p> <p>MT group: 13 patients. Men: 1 Women: 12 Age Range: 20.9–61.5</p> <p>Combined</p>	None.

		therapies (CTs) group: 14 patients. Men:1 Women: 13 Age range: 32.1– 61.9.	
--	--	---	--

3.3. Study Characteristics

[Table 4](#) shows the main characteristics of the analyzed studies with the most important information being narrowed down to clarify the hypotheses proposed by this study. The type of study, the sample size, the intervention performed, the variables analyzed, and the results obtained, with their respective numerical values, are among the properties found.

3.4. Methodological Quality of the Collected

Literature

Two scales were used to carry out the methodological quality of the present review. The PEDro scale resulted in eleven articles being selected of which six were given an "excellent" score and the remaining five a "good" score. The IVS resulted in seven articles receiving a "high" score and four were "limited".

3.5. Effects of Photobiomodulation Therapy on Health Improvement and Chronic Pain

The articles in this study were chosen in order to determine the efficacy of PBMT application on subjects with health disorders such as pain or inflammation.

In the subsequent points, the results of the highlighted variables from the selected studies are presented.

3.5.1. Chronic Pain

In this study, 11 articles have been selected that deal with different types of chronic pain.

In the studies by Fernanda R. et al. [18], Leyla K. et al. [19], Mariana M. et al. [20], De Souza R. et al. [21], Shaiane S. et al. [22], Taradaj J. et al. [24], and Fang-Y L. et al. [25] PBMT was applied to patients with chronic pain⁴

Two of these studies, by Fernanda R, et al. [18] and Fang-Y L. et al. [25], deal with female patients, over the age of 60, with knee osteoarthritis. Both studies revealed significant improvements in the patients, with p -values less than 0.05 in all the studied variables related to chronic pain.

On the other hand, we find the studies by Leyla K. et al. [19], Shaiane S. et al. [22], and Taradaj J. et al. [24] which have subjects with nonspecific chronic low back pain. In the first of these, in which there was an equal ratio of men and women and an average age of 49, LLLT produced significant changes in pain and functional status ($p < 0.001$) during the 3-month treatment period, unlike the placebo group that only saw these changes for the first month. The study by Taradaj J. et al. [24] also included subjects of both sexes with an average age of 45. The results showed that the changes produced by PBMT were short-term (3 weeks after therapy), since the 1- and 3-month evaluations resulted in a $p > 0.05$ when the exercises were not continued. The last remaining study is that of Shaiane S. et al. [22], which has a mean age of 34 and includes both male and female participants. The study indicates that PGE2 significantly reduced ($p = 0.04$), therefore, the pain decreased.

The studies by Mariana M. et al. [20] y De Souza R. et al. [21] are comprised of participants with fibromyalgia, and also focus on pain, among other variables. In the first study, the

participants are exclusively female, while in the second only the majority are women. The average age of the first study is 35, with 46 being the average age in the second. The study by Mariana M. et al. [20] found that a PBMT session was markedly better than an exercise session, however, no additional benefits were found with a combination of both therapies. In the long term (10 weeks) the combined therapy group and the PBMT group experienced a significant reduction in pain compared to the placebo and the therapeutic exercise groups. The De Souza R. et al. [21] study evaluated the orofacial pain of these FM patients, and the results indicate a significant decrease in pain ($p = 0.0001$) due to the treatment, although with no noticeable difference between the two groups.

3.5.2. Inflammation

In the studies by Gonçalves Langella L. et al. [17], Shaiane S. et al. [22], and Chong R. et al. [19] the fundamental part of the research was to observe the effect of PBMT on the most important biomarkers related to inflammation. The first of these studies observed changes in the serum levels of the proinflammatory cytokines IL-6, IL-8, and TNF- α , in 18 patients after knee arthroplasty surgery, pointing to the fact that PBMT significantly reduced inflammation ($p < 0.05$) when compared to the placebo group. The second of these studies also measured the serum levels of IL-6 and TNF- α in

18 patients with nonspecific chronic low back pain after applying LLLT, but unlike the first, the serum changes in these cytokines did not cause a significant reduction in pain. The last of these studies looked at lipid biomarkers associated with pain and inflammation in 54 people with orthodontic pain and periodontal inflammation. It was noted that the LLLT group had significant improvements in pain 1 h and 24 h later ($p = 0.02$) and in inflammation during the first 24 h.

3.5.3. Acute Pain

Nunes EC. et al. [20] and Gonçalves Langella L. et al. [17] tried to observe how PBMT affects pain but in this case acute pain. The former study, in which groups with a similar male/female ratio participated, used the VPRS and the NPRS since the participants were post-endodontic surgery patients. The study showed that the reduction in pain with laser therapy was significant ($p < 0.001$). The second study also indicates a significant reduction in pain, but this time after knee arthroplasty surgery ($p < 0.05$).

Fernanda T. et al. [26] evaluated temporomandibular pain in a study made up of 41 patients with temporomandibular dysfunction and pain. The groups received PBMT, manual therapy, or a combination of both. The results revealed that there were significant changes in pain reduction ($p < 0.001$), with all managing to reduce it significantly, but this

difference did not exist between groups.

3.6. Questionnaires and Scales Used in the Articles

A visual analog scale (VAS) is used in the quantitative measurement of pain. It is a linear measurement in which the ends reflect the minimum level at one end and the maximum level of pain at the other. The scale contains a single item that can give a score from 0 to 10. Normally, "no pain" results in the number 0, and "very intense pain/worst imaginable pain " results in 10 [27].

The Fibromyalgia Impact Questionnaire (FIQ) consists of a three-factor structure with a physical symptoms domain (six items), a functional domain (ten items), and a mental symptoms domain (two items) [28].

Research Diagnostic Criteria (RDC) classify subjects with temporomandibular disorders following a physical diagnosis and their status related to pain disability and psychological functioning [29].

The Roland Morris Questionnaire (RMQ) measures low back pain, consisting of an action list (24 items) that usually affects low back pain [30,31].

Numerical Pain Scale (NPRS): It was developed with the

purpose of unidimensionally assessing adult pain. It is an alternative version of the VAS, using a horizontal line, with 11 items, and a score range of 0–10, with 0 being “no pain” and 10 being “more intense pain than can be imagined” [27].

Western Ontario and McMaster Universities Osteoarthritis Index (WOMAC) measures and evaluates pain, functional capacity, and stiffness in osteoarthritic knee and hip diseases. It is made up of 24 items that are divided into three scales. Each item is answered using a verbal scale which has five levels (0–4) [12].

The Beck Anxiety Inventory (BAI): This inventory was designed to assess anxiety in the cognitive and physical domains in the last week before completing the questionnaire. It consists of 21 items and the score can be between 0 and 63. Levels 0 to 7 mean minimal anxiety, 8 to 15 mean mild anxiety, 16 to 25 mean moderate anxiety 16–25, and 26 to 63 mean severe anxiety [32].

4. Discussion

The goal of this review was to evaluate the current knowledge on the efficacy of PBM in chronic musculoskeletal pain, acute pain, and inflammation treatment plans. Moreover, to assess the methodological quality of the included studies.

The search and selection process resulted in 11 articles which were all randomized clinical trials.

Of the selected trials, seven dealt with how chronic pain is or is not modified after the application of PBM [16,18,19,20,21,23,33,34], with the VAS being used as the main element for measuring pain.

There are articles [17,18,19] that focus on the impact PBM has on inflammation. The first two studies check its efficacy by measuring the serum levels of the proinflammatory cytokines IL-6, IL-8, and TNF- α . The latter study [19] observed the FGC biomarkers associated with inflammation.

Among the selected studies, there are several that use the VAS to quantitatively measure the changes, and also study the effect of low-intensity laser on patients with acute pain [16,19,20].

In the following sections, we will analyze the impact that PBM has had on different conditions.

4.1. Chronic Nonspecific Low Back Pain

Chronic nonspecific low back pain is the most prevalent painful musculoskeletal condition. It is heterogeneous since it can have many causes and diagnoses, but there are very few therapies with solid evidence of efficacy [33].

Low-level light therapy has played a different role in each study. In one study [33] the EG consisted of seven men and thirteen women with an average age of 47 years. After the interventions were carried out and the results evaluated, the subjects were found to have experienced significant improvements in pain and functional state with the improvements lasting for the 3-month treatment period. On the other hand, a study [23], with an average age of 34 years and made up of four men and five women did not find a significant improvement in pain, unlike in the aforementioned study. The methods and parameters of the therapy application were similar, although not identical. In the last of the studies the average age of the sixteen subjects in the EG, eight men and eight women was 45.5. This study concluded that the application of PBM had a significant influence on mobility improvement and hence pain, but only in the short term; the parameters deteriorated at the 1- and 3-month post-therapy analyses where there were no exercises [19].

Knee osteoarthritis

Osteoarthrosis, a very prevalent musculoskeletal condition, is thought to affect 10% of men and 13% of women who are 60 years of age or older. This pathology affects the majority of the affected joints' tissues and structures, which produces structural and functional changes [3].

The first of the selected studies was made up of two experimental groups, one received US + LLLT while the other received the same with the addition of ET. The age range of the women who made up each group ranged from 61 to 77 years. The results showed a significant increase in pain threshold, and the number of squats in the experimental group when compared to the control group.

Another RCT on PBM was conducted in patients with osteoarthritis. There were 16 patients of both sexes between the ages of 63 and 78 who received PBM three times per week for 4 weeks. Mobility was significantly improved each week compared to the CG. The same occurred with pain threshold and the knee OA severity index [16].

4.2. Fibromyalgia

This syndrome is common and is characterized by widespread chronic pain, fatigue, and sleep disorders that severely attack those affected. The etiopathogenesis of this syndrome is not clear, but therapy is available to improve quality of life [35].

In the first of the selected studies with patients suffering from FM, two sets were formed that received the same therapies, namely PBM, ET, and PBM + ET. Set 1 was used to evaluate the therapies in the short term and set 2 in the long term. The age range of 32–42 years old in the two sets was

similar. The results determined that the ET group was the only group in set 1 not to significantly improve pain threshold. The PBM + ET group showed no additional benefits in set 1 but did in set 2, which examined the changes over a 10-week period, thus showing that PBM and PBM + ET produced significant improvements in pain threshold. Both combined therapies reduced the number of tender points and had a beneficial role in patients' anxiety, depression, and fatigue [20].

The second study, unlike the previous one, did not have a CG, but instead had two EGs each comprised of 33 people suffering from FM and with an average age of 46 years. One group received PBM and the other an aesthetic infiltration of lidocaine. The group that received PBM showed a significant decrease in the intensity of orofacial pain, in the same way as the group that received the infiltration. In both groups, all the analyzed study variables improved significantly, although the patients' perception of well-being was better for the group that received laser irradiation [21].

Current research shows promising short-term results of whole-body PBM on fibromyalgia sufferers, however, the long-term results are still to be explored [6,7].

4.3. Temporomandibular Pain and Dysfunction

This disorder is a very common condition experienced

mainly by young middle-aged women. The etiology is unknown but the following risk factors seem to be associated with it: bruxism, depression, multiple pain conditions, gender, trauma, and hypermobility [36].

In this study, we incorporated an RCT that verifies the efficacy of photobiomodulation in this condition. It was verified by applying PBM three times a week for a month. The study had no CG but had three EGs receiving, TFBM, TM, or a combination of both. The study had 41 subjects, only 2 of them men. All groups experienced a significant reduction in pain and anxiety. PBM promoted improvements in five conditions, TM in two, and combined therapies in one [17].

4.4. Oral Pain and Inflammation

Two selected RCTs tried to report on the impact of PBM on patients with periodontal and endodontic pain and inflammation. The first applied laser therapy to a group of 35 people ranging from 20–37 years old with similar male/female ratios. This article highlights the significant improvement that the therapy produced during the first 24 h when compared to the placebo group [25].

The other group had an EG of 27 people who, after receiving the interventions, were evaluated for subjective pain and the FGC biomarkers associated with pain. The group that

received the PBM obtained improvements 1 h and 24 h after treatment. Additionally, subjective pain was also significantly better than in the CG at an interval of 3 days post-application [24].

4.5. Post-Surgery Pain and Swelling

Inflammation is known to be the body's response to an injury to our system. Inflammatory mediators induce pain through the direct activation of nociceptors, hence the two are closely related concepts [37].

One article reviewed the influence of PBM on acute pain and swelling after knee arthroplasty. To accomplish this, nine patients in the EG had an application of PBM. The results revealed a positive and significant improvement in both pain and inflammation post-therapy on five points along the surgical scar [22].

4.6. Strengths and Weaknesses of the Study

The present review shows the following strengths:

Good methodological quality and therefore the results are assured by the use of the PEDro scale.

The wide variety of medical conditions presented by the subjects of the selected studies.

The study highlights the lack of studies on this subject and hence the proposal for a novel therapy that can non-invasively address chronic musculoskeletal pain and inflammation.

As weak points there are certain limitations:

Limited RCTs showing the influence of PBM on subjects with inflammation.

A lack of studies that describe the ideal standard parameters according to the type of condition. Since only English studies were included, the extent of the study is reduced.

Heterogeneity of the study subjects, due to the fact that chronic musculoskeletal pain conditions come from different etiologies. Therefore, biomarkers and pain-generating mechanisms may differ.

4.7. Prospective

After observing and analyzing the results, it can be seen that the scientific evidence on PBM in relation to chronic pain and inflammation is scarce. Therefore, different lines of research are proposed as follows:

Implementation of RCTs on the efficacy of PBM application strategies in subjects with different kinds of inflammation.

Conduct RCTs on the efficacy of PBM application strategies in subjects with chronic musculoskeletal pain.

Perform RCTs that try to discern standard parameters according to the type of pain or inflammation, in order to move towards a consensus and be able to research the influence of laser therapy under standardized conditions.

Although new evidence analyzing the effects of PBM on sports performance has been carried out [38], studies that show the effects of PBM on elite athletes suffering from pain are still needed.

Finally, studies in subjects below 18 years of age who suffer from acute or chronic pain are necessary.

5. Conclusions

The quality of the studies collected on the application of PBM in the treatment of inflammation and chronic pain is generally high, so they have excellent quality in terms of methodology.

The symptoms of pain and chronic pain are influenced by laser irradiation. However, there is a lack of studies to clarify this ability to reduce pain, especially in the long term.

There is not enough evidence on the effects and benefits of PBM on inflammation. Although the studies confirm the

positive effects that this therapy has on proinflammatory biomarkers, there are only a few RCTs that verify this, and more research is needed.

Funding Statement

This study has been partially funded by the University Chair in Clinical Psychoneuroimmunology (University of Granada and PNI Europe).

Author Contributions

Study conception (S.N.-L. and A.G.-M.), design (S.N.-L. and A.G.-M.), acquisition of data (A.G.-M.), analysis and interpretation of data (S.N.-L., A.G.-M., M.C.-C., M.A.-G., D.H.-H., J.J.P.-M. and D.A.-N.) drafting of manuscript (S.N.-L., A.G.-M., M.C.-C., M.A.-G., L.P., D.H.-H., J.J.P.-M. and D.A.-N.) and critical revision (S.N.-L., A.G.-M., M.C.-C., M.A.-G., D.H.-H., J.J.P.-M., L.P. and D.A.-N.). All authors have read and agreed to the published version of the manuscript.

Institutional Review Board Statement

Not applicable.

Informed Consent Statement

Not applicable.

Data Availability Statement

All data associated with this study are present in the paper. All requests for other materials will be reviewed by the corresponding author to verify whether the request is subject to any intellectual property or confidentiality obligations.

Conflicts of Interest

The authors declare no conflict of interest.

Footnotes

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.

References

1. Salehpour F., Mahmoudi J., Kamari F., Sadigh-Eteghad S., Rasta S.H., Hamblin M.R. Brain Photobiomodulation Therapy: A Narrative Review. *Mol. Neurobiol.* 2018;55:6601–6636. doi: 10.1007/s12035-017-0852-4. [[PMC free article](#)]

[\[PubMed\]](#) [\[CrossRef\]](#) [\[Google Scholar\]](#)

2. Karu T.I., Pyatibrat L.V., Kolyakov S.F., Afanasyeva N.I. Absorption measurements of a cell monolayer relevant to phototherapy: Reduction of cytochrome c oxidase under near IR radiation. *J. Photochem. Photobiol. B Biol.* 2005;81:98–106. doi: 10.1016/j.jphotobiol.2005.07.002.

[\[PubMed\]](#) [\[CrossRef\]](#) [\[Google Scholar\]](#)

3. Yeh S.W., Hong C.H., Shih M.C., Tam K.W., Huang Y.H., Kuan Y.C. Low-level laser therapy for fibromyalgia: A systematic review and meta-analysis. *Pain Physician.* 2019;22:241–254. [\[PubMed\]](#) [\[Google Scholar\]](#)

4. Gur A., Karakoc M., Cevik R., Nas K., Sarac A.J., Karakoc M. Efficacy of low power laser therapy and exercise on pain and functions in chronic low back pain. *Lasers Surg. Med.* 2003;32:233–238. doi: 10.1002/lsm.10134. [\[PubMed\]](#) [\[CrossRef\]](#) [\[Google Scholar\]](#)

5. Honda Y., Sakamoto J., Hamaue Y., Kataoka H., Kondo Y., Sasabe R., Goto K., Fukushima T., Oga S., Sasaki R., et al. Effects of Physical-Agent Pain Relief Modalities for Fibromyalgia Patients: A Systematic Review and Meta-Analysis of Randomized Controlled Trials. *Pain Res. Manag.* 2018;2018:1–9. doi: 10.1155/2018/2930632. [\[PMC free article\]](#) [\[PubMed\]](#) [\[CrossRef\]](#) [\[Google Scholar\]](#)

6. Navarro-Ledesma S., Carroll J., Burton P., Ana G.-M. Short-Term Effects of Whole-Body Photobiomodulation on Pain, Quality of Life and Psychological Factors in a Population Suffering from Fibromyalgia: A Triple-Blinded

Randomised Clinical Trial. *Pain Ther.* 2022;12:225–239.

doi: 10.1007/s40122-022-00450-5. [[PMC free article](#)]

[[PubMed](#)] [[CrossRef](#)] [[Google Scholar](#)]

7. Navarro-Ledesma S., Carroll J., González-Muñoz A., Pruijboom L., Burton P. Changes in Circadian Variations in Blood Pressure, Pain Pressure Threshold and the Elasticity of Tissue after a Whole-Body Photobiomodulation Treatment in Patients with Fibromyalgia: A Triple-Blinded Randomized Clinical Trial. *Biomedicines.* 2022;10:2678.

doi: 10.3390/biomedicines10112678. [[PMC free article](#)]

[[PubMed](#)] [[CrossRef](#)] [[Google Scholar](#)]

8. Wickenheisser V.A., Zygot E.M., Rabjohns E.M., Lee H.H., Lawrence D.S., Tarrant T.K. Laser Light Therapy in Inflammatory, Musculoskeletal, and Autoimmune Disease.

Curr. Allergy Asthma Rep. 2019;19:37. doi: 10.1007/s11882-

019-0869-z. [[PMC free article](#)] [[PubMed](#)] [[CrossRef](#)]

[[Google Scholar](#)]

10. Mills S.E.E., Nicolson K.P., Smith B.H. Chronic pain: A review of its epidemiology and associated factors in population-based studies. *Br. J. Anaesth.* 2019;123:e273–

e283. doi: 10.1016/j.bja.2019.03.023. [[PMC free article](#)]

[[PubMed](#)] [[CrossRef](#)] [[Google Scholar](#)]

11. Domenichiello A.F., Ramsden C.E. The silent epidemic of chronic pain in older adults. *Prog. Neuro-Psychopharmacol. Biol. Psychiatry.* 2019;93:284–290.

doi: 10.1016/j.pnpbp.2019.04.006. [[PMC free article](#)]

[[PubMed](#)] [[CrossRef](#)] [[Google Scholar](#)]

12. Kaushik A.S., Strath L.J., Sorge R.E. Dietary Interventions for Treatment of Chronic Pain: Oxidative Stress and Inflammation. *Pain Ther.* 2020;9:487–498.

doi: 10.1007/s40122-020-00200-5. [[PMC free article](#)]

[[PubMed](#)] [[CrossRef](#)] [[Google Scholar](#)]

13. Bosch-Capblanch X., Lavis J.N., Lewin S., Atun R., Røttingen J.-A., Dröschel D., Beck L., Abalos E., El-Jardali F., Gilson L., et al. Policy Forum Guidance for Evidence-Informed Policies about Health Systems: Rationale for and Challenges of Guidance Development. *PLoS Med.*

2012;9:e1001185. doi: 10.1371/journal.pmed.1001185. [[PMC](#)

[free article](#)] [[PubMed](#)] [[CrossRef](#)] [[Google Scholar](#)]

14. Clijsen R., Brunner A., Barbero M., Clarys P., Taeymans J. Effects of low-level laser therapy on pain in patients with musculoskeletal disorders: A systematic review and meta-analysis. *Eur. J. Phys. Rehabil. Med.* 2017;53:603–610.

doi: 10.23736/S1973-9087.17.04432-X. [[PubMed](#)]

[[CrossRef](#)] [[Google Scholar](#)]

15. Maher C.G., Sherrington C., Herbert R.D., Moseley A.M., Elkins M. Reliability of the PEDro Scale for Rating Quality of Randomized Controlled Trials. *Phys. Ther.* 2003;83:713–721.

doi: 10.1093/ptj/83.8.713. [[PubMed](#)] [[CrossRef](#)] [[Google](#)

[Scholar](#)]

16. Nunes E.C., Herkrath F.J., Suzuki E.H., Júnior E.C.G., Marques A.A.F., Júnior E.C.S. Comparison of the effect of photobiomodulation therapy and Ibuprofen on postoperative pain after endodontic treatment: Randomized, controlled,

clinical study. *Lasers Med Sci.* 2019;35:971–978.

doi: 10.1007/s10103-019-02929-8. [[PubMed](#)] [[CrossRef](#)]
[[Google Scholar](#)]

17. Langella L.G., Casalechi H.L., Tomazoni S.S., Johnson D.S., Albertini R., Pallotta R.C., Marcos R.L., Carvalho P.D.T.C.D., Leal-Junior E.C.P. Photobiomodulation therapy (PBMT) on acute pain and inflammation in patients who underwent total hip arthroplasty—a randomized, triple-blind, placebo-controlled clinical trial. *Lasers Med Sci.*

2018;33:1933–1940. doi: 10.1007/s10103-018-2558-x.

[[PubMed](#)] [[CrossRef](#)] [[Google Scholar](#)]

18. Paolillo F.R., Paolillo A.R., João J.P., Frascá D., Duchêne M., João H.A., Bagnato V.S. Ultrasound plus low-level laser therapy for knee osteoarthritis rehabilitation: A randomized, placebo-controlled trial. *Rheumatol. Int.* 2018;38:785–793.

doi: 10.1007/s00296-018-4000-x. [[PubMed](#)] [[CrossRef](#)]

[[Google Scholar](#)]

19. Kholoosy L., Elyaspour D., Akhgari M.R., Razzaghi Z., Khodamardi Z., Bayat M. Evaluation of the therapeutic effect of low level laser in controlling low back pain: A randomized controlled trial. *J. Lasers Med. Sci.* 2020;11:120–125.

doi: 10.34172/jlms.2020.21. [[PMC free article](#)] [[PubMed](#)]

[[CrossRef](#)] [[Google Scholar](#)]

20. da Silva M.M., Albertini R., Carvalho P.D.T.C.D., Leal-Junior E.C.P., Bussadori S.K., Vieira S.S., Bocalini D.S., de Oliveira L.V.F., Grandinetti V., Silva J.A., et al. Randomized, blinded, controlled trial on effectiveness of

photobiomodulation therapy and exercise training in the fibromyalgia treatment. *Lasers Med Sci.* 2017;33:343–351. doi: 10.1007/s10103-017-2388-2. [[PubMed](#)] [[CrossRef](#)] [[Google Scholar](#)]

21. de Souza R.-C.-V., de Sousa E.-T., Oliveira Scudine K.-G., Meira U.-M., de Oliveira Silva E.-D., Gomes A.-C.-A., Limeira-Junior F.-d.A. Low-level laser therapy and anesthetic infiltration for orofacial pain in patients with fibromyalgia: A randomized clinical trial. *Med. Oral Patol. Oral Y Cir. Bucal.* 2018;23:e65–e71. doi: 10.4317/medoral.21965. [[PMC free article](#)] [[PubMed](#)] [[CrossRef](#)] [[Google Scholar](#)]

22. Tomazoni S.S., Costa L.O.P., Joensen J., Stausholm M.B., Naterstad I.F., Ernberg M., Leal-Junior E.C.P., Bjordal J.M. Photobiomodulation Therapy is Able to Modulate PGE 2 Levels in Patients with Chronic Non-Specific Low Back Pain: A Randomized Placebo-Controlled Trial. *Lasers Surg. Med.* 2020;53:236–244. doi: 10.1002/lsm.23255. [[PubMed](#)] [[CrossRef](#)] [[Google Scholar](#)]

23. Ren C., McGrath C., Gu M., Jin L., Zhang C., Sum F.H.K.M.H., Wong K.W.F., Chau A.C.M., Yang Y. Low-level laser-aided orthodontic treatment of periodontally compromised patients: A randomised controlled trial. *Lasers Med Sci.* 2019;35:729–739. doi: 10.1007/s10103-019-02923-0. [[PubMed](#)] [[CrossRef](#)] [[Google Scholar](#)]

24. Taradaj J., Rajfur K., Rajfur J., Ptaszkowski K., Ptaszkowska L., Sopol M., Rosińczuk J., Dymarek R. Effect of laser treatment on postural control parameters in patients

with chronic nonspecific low back pain: A randomized placebo-controlled trial. *Braz. J. Med. Biol. Res.* 2019;52 doi: 10.1590/1414-431x20198474. [[PMC free article](#)]

[[PubMed](#)] [[CrossRef](#)] [[Google Scholar](#)]

25. Liao F.-Y., Lin C.-L., Lo S.-F., Chang C.-C., Liao W.-Y., Chou L.-W. Efficacy of Acupoints Dual-Frequency Low-Level Laser Therapy on Knee Osteoarthritis. *Evid.-Based Complement. Altern. Med.* 2020;2020:1–7.

doi: 10.1155/2020/3951741. [[PMC free article](#)] [[PubMed](#)]

[[CrossRef](#)] [[Google Scholar](#)]

26. Brochado F.T., De Jesus L.H., Carrard V.C., Freddo A.L., Chaves K.D., Martins M.D. Comparative effectiveness of photobiomodulation and manual therapy alone or combined in TMD patients: A randomized clinical trial. *Braz. Oral Res.* 2018;32:e50. doi: 10.1590/1807-3107bor-2018.vol32.0050.

[[PubMed](#)] [[CrossRef](#)] [[Google Scholar](#)]

[[PubMed](#)] [[CrossRef](#)] [[Google Scholar](#)]

27. Sung Y.-T., Wu J.-S. The Visual Analogue Scale for Rating, Ranking and Paired-Comparison (VAS-RRP): A new technique for psychological measurement. *Behav. Res. Methods.* 2018;50:1694–1715. doi: 10.3758/s13428-018-

1041-8. [[PMC free article](#)] [[PubMed](#)] [[CrossRef](#)] [[Google](#)

[Scholar](#)]

[[PubMed](#)] [[CrossRef](#)] [[Google Scholar](#)]

28. Van Wilgen C.P., Vuijk P., Van Ittersum M.W., Nijs J. Not throwing out the baby with the bathwater: Lessons from the Fibromyalgia Impact Questionnaire. *Clin. Rheumatol.* 2012;32:333–339. doi: 10.1007/s10067-012-2120-3.

[[PubMed](#)] [[CrossRef](#)] [[Google Scholar](#)]

[[PubMed](#)] [[CrossRef](#)] [[Google Scholar](#)]

29. Winocur E., Steinkeller-Dekel M., Reiter S., Eli I. A retrospective analysis of temporomandibular findings among Israeli-born patients based on the RDC/TMD. *J. Oral Rehabil.* 2009;36:11–17. doi: 10.1111/j.1365-2842.2008.01898.x. [[PubMed](#)] [[CrossRef](#)] [[Google Scholar](#)]
30. Madan J., Khan K.A., Petrou S., Lamb S.E. Can Mapping Algorithms Based on Raw Scores Overestimate QALYs Gained by Treatment? A Comparison of Mappings between the Roland-Morris Disability Questionnaire and the EQ-5D-3L Based on Raw and Differenced Score Data. *Pharmacoeconomics.* 2017;35:549–559. doi: 10.1007/s40273-016-0483-z. [[PubMed](#)] [[CrossRef](#)] [[Google Scholar](#)]
31. Primorac D., Molnar V., Rod E., Jeleč Ž., Čukelj F., Matišić V., Vrdoljak T., Hudetz D., Hajsok H., Borić I. Knee Osteoarthritis: A Review of Pathogenesis and State-Of-The-Art Non-Operative Therapeutic Considerations. *Genes.* 2020;11:854. doi: 10.3390/genes11080854. [[PMC free article](#)] [[PubMed](#)] [[CrossRef](#)] [[Google Scholar](#)]
32. Oehlert M.E., Nelson K.G., King N., Reis D.J., Sumerall S., Neal C., Henry P. Measurement-based care: Use of the beck anxiety inventory (BAI) in a veteran population. *Psychol. Serv.* 2020;17:372–379. doi: 10.1037/ser0000372. [[PubMed](#)] [[CrossRef](#)] [[Google Scholar](#)]
33. Oliveira I.S., Tomazoni S.S., Vanin A.A., Araujo A.C., De Medeiros F.C., Oshima R.K.A., Costa L.O.P., Costa L.D.C.M. Management of Acute Low Back Pain in Emergency

Departments in São Paulo, Brazil: A Descriptive, Cross-Sectional Analysis of Baseline Data from a Prospective Cohort Study. *BMJ Open*. 2022;12 doi: 10.1136/bmjopen-2021-059605. [[PMC free article](#)] [[PubMed](#)] [[CrossRef](#)] [[Google Scholar](#)]

34. Thomsen M.G., Latifi R., Kallemose T., Barfod K.W., Husted H., Troelsen A. Good validity and reliability of the forgotten joint score in evaluating the outcome of total knee arthroplasty. *Acta Orthop*. 2016;87:280–285.

doi: 10.3109/17453674.2016.1156934. [[PMC free article](#)] [[PubMed](#)] [[CrossRef](#)] [[Google Scholar](#)]

35. Navarro-Ledesma S., Gonzalez-Muñoz A., Carroll J., Burton P. Short- and Long-Term Effects of Whole-Body Photobiomodulation on Pain, Functionality, Tissue Quality, Central Sensitisation and Psychological Factors in a Population Suffering from Fibromyalgia: Protocol for a Triple-Blinded Randomised Clinical Trial. *Ther. Adv. Chronic. Dis*. 2022;13:204062232210780.

doi: 10.1177/20406223221078095. [[PMC free article](#)] [[PubMed](#)] [[CrossRef](#)] [[Google Scholar](#)]

38. González-Muñoz A., Perez-Montilla J.J., Cuevas-Cervera M., Aguilar-García M., Aguilar-Nuñez D., Hamed-Hamed D., Pruiomboom L., Navarro-Ledesma S. Effects of Photobiomodulation in Sports Performance: A Literature Review. *Appl. Sci*. 2023;13:3147. doi: 10.3390/app13053147. [[CrossRef](#)] [[Google Scholar](#)]

Articles from Healthcare are provided here courtesy of
Multidisciplinary Digital Publishing Institute (MDPI)