

# **Brain Waste Removal System and Sleep: Photobiomodulation as an Innovative Strategy for Night Therapy of Brain Diseases**

## **Abstract**

Emerging evidence suggests that an important function of the sleeping brain is the removal of wastes and toxins from the central nervous system (CNS) due to the activation of the brain waste removal system (BWRS). The meningeal lymphatic vessels (MLVs) are an important part of the BWRS. A decrease in MLV function is associated with Alzheimer's and Parkinson's diseases, intracranial hemorrhages, brain tumors and trauma. Since the BWRS is activated during sleep, a new idea is now being actively discussed in the scientific community: night stimulation of the BWRS might be an innovative and promising strategy for neurorehabilitation medicine. This review highlights new trends in photobiomodulation of the BWRS/MLVs during deep sleep as a breakthrough technology for the effective removal of wastes and unnecessary compounds from the brain in order to increase the neuroprotection of the CNS as

well as to prevent or delay various brain diseases.

**Keywords:** brain diseases, meningeal lymphatic vessels, photobiomodulation

# **1. Therapeutic Properties of Activation of Brain Waste Removal System (BWRS) during Deep Sleep**

What do we sleep for? A widespread belief is that sleep is a crucial function of the brain that is necessary for our recharging, leaving our energy refreshed when we wake up. Good sleep also helps to maintain our health and protects against the development of various diseases, including brain pathologies [1]. When we do not get enough sleep, our brain and body cannot function properly. Long-term sleep deficit can lead to the formation of dementia. Indeed, the results of a recent study on a large group of 8000 volunteers observed from their middle age till their 70th year of age for 25 years revealed a higher incidence of dementia in people aged 50–60 years who chronically did not get enough sleep (6 h and less a night) [2]. It is interesting to note that even one night of sleep deprivation causes an increase in the level of beta-amyloid ( $A\beta$ ) protein in the brain of young and healthy volunteers [3,4]. There is a growing body of evidence suggesting that sleep disturbance is an independent risk factor of cognitive impairment, and the measurement of sleep quality can be an innovative method to screen for

Alzheimer's disease (AD) [5,6,7,8,9,10,11,12,13,14]. It is well known that people with AD experience poor and short sleep, which is associated with increased A $\beta$  deposition in their brains [8,11]. Experimental and clinical studies have shown that the A $\beta$  content in cerebral spinal fluid (CSF) is the highest in the evening before sleep and the lowest in the morning after sleep [3,15]. There is the hypothesis that sleep is accompanied by A $\beta$  clearance from the brain [3,15]. It was recently discovered that sleep deficit causes the opening of the blood–brain barrier (BBB) to inflammatory mediators and immune cells in both humans and rodents [16,17,18,19,20]. Sleep is considered an important biomarker and a promising therapeutic target for cerebral small-vessel diseases, including AD and brain pathologies associated with BBB disruption [16]. Chronic sleep restriction promotes astrocytic phagocytosis of synaptic elements and microglia activation, i.e., the brain begins to “eat” itself [21]. Obviously, sleep is essential for the health of the central nervous system (CNS). However, the mechanisms underlying the phenomenon of restorative sleep remain unknown.

Aristotle came up with the brilliant idea that sleep triggers brain cleansing [22]. After 2000 years, his idea was confirmed by several studies suggesting that sleep activates unique brain processes for the removal of metabolites and wastes [15,23,24,25,26,27,28] (Figure 1). However, not all sleep is crucial to CNS health. There are two types of sleep:

non-rapid eye movement (NREM) or deep sleep and rapid eye movement (REM) sleep, when we dream. Current studies demonstrate that only NREM sleep is optimal for functions of the BWRS [29,30]. BWRS activity is significantly increased during NREM sleep and is dramatically suppressed during wakefulness [15]. Animal data clearly demonstrate a 95% reduction in A $\beta$  removal from the brain during wakefulness and the activation of this process during NREM sleep due to a 60% increase in interstitial fluid (ISF) space [15]. Later, it was confirmed that sleep induces the enhancement of BWRS activity arising from the expansion of ISF space [31]. Body posture during sleep is also important for the optimization of BWRS function [32]. Lee et al. reported that the horizontal position of the body during sleep contributes the most to optimal waste removal, including A $\beta$  clearance from the brain [32].

**Figure 1.**

[Open in a new tab](#)

BWRS activation during sleep. During wakefulness, the BWRS is inactive; therefore, wastes and toxins accumulate in the brain. During deep sleep, the BWRS is activated, which is

accompanied by an increase in the drainage of brain tissues and the intense elimination of wastes and toxins. Figuratively speaking, the brain, during deep sleep, turns into a washing machine, removing unnecessary compounds from its tissues.

The BWRS is the most efficient during slow-wave activity (SWA; 0–4 Hz), which is a major rhythm and a marker of NREM [29,30]. The physiological significance of SWA remains a mystery, but there is growing evidence that SWA plays an essential role in the control of sleep quality and the optimization of BWRS function [5,15,30,33,34,35,36,37]. SWA is strongly controlled, and sleep loss immediately causes a compensatory increase in SWA time during subsequent nocturnal sleep [33]. Interestingly, it is possible to suppress REM sleep in mice [38]. However, among the hundreds of mutant mouse models, no mice have yet been discovered that exhibit NREM sleep loss [39]. SWA is considered brain activity supporting the sleep regulation of the BWRS [23]. Results of contrast-enhanced magnetic resonance imaging (MRI) reveal that the circadian rhythm is an important driving force for the movement of wastes in ISF space and their clearance from the brain [40,41]. The deletion of aquaporin channels (AQP4) eliminates the effects of the circadian rhythm on brain fluid transport and the clearance of proteins from the brain [41]. Since AQP4 in astrocytic endfeet is controlled by the circadian rhythm, AQP4 optimizes brain fluid movement and waste clearance [29,41,42]. There is evidence that astrocytes control circadian timekeeping via glutamate signaling [43]. Thus,

astrocytes and AQP4 present a checkpoint for BWRS function during NREM sleep [29,44]. These pioneering findings were obtained in rodents, whose circadian rhythm differs significantly from that of humans [45]. Presently, the exact contribution of the circadian rhythm to the optimization of processes of waste clearance from the human brain remains unknown [41]. Therefore, further studies are necessary to confirm the circadian control of the BWRS in humans. In addition, SWA is the result of metabolic and neurochemical changes in the sleeping brain that provide the predominance of specific neural network oscillations associated with changes in different physiological parameters, such as blood pressure, heart rate and respiratory changes, which can also regulate the BWRS [46,47,48,49].

Another factor that concerns SWA is temperature. During NREM sleep, the mouse brain neocortex temperature decreases by  $\sim 2$  °C [50,51]. During this time, the hypothalamus stimulates the cooling of the brain, which induces NREM sleep [52]. There is the hypothesis that the cooling of the brain is a key function of sleep to preserve energy that might be linked to synaptic remodeling associated with SWA [53]. Cooling during NREM sleep stimulates the expression of the cold-inducible RNA-binding protein and RNA-binding motif protein 3 genes that are required for structural remodeling (they are also induced in

some hibernators) [53]. The “cooling and cleaning” of the brain is a physiological phenomenon and an informative platform for therapy in degenerative and neuro-inflammatory diseases [1,54].

Taken together, considerable evidence suggests a functional relationship between sleep and the BWRS. The BWRS is considered a brain function related to waste and fluid transport with astrocyte-regulated mechanisms, while BWRS dysfunction is associated with different brain diseases, especially with cognitive decline [9,15,29,37,55,56,57,58,59,60,61,62,63]. Indeed, SWA is a new therapeutic target for AD [9,37]. Patients and animals with AD demonstrate SWA disturbances [11,37,64,65,66]. In humans,  $A\beta$  accumulation in the brain is correlated with both decreased SWA time and impaired memory consolidation [67,68]. SWA disruption is also reported in patients with mild cognitive impairment [69]. It is assumed that sleep impairment in cognitively normal old people could predict  $A\beta$  and tau accumulation in the brain [70]. Why SWA reflects AD pathology remains unknown. Decreased BWRS function after insufficient sleep may be related to an increase in the interstitial noradrenaline (NE) level [15], leading to a reduction in astrocytic volume [71] and the vasoconstriction of the pial arteries [72].  $A\beta$ -mediated abnormalities in excitatory and inhibitory neurotransmission can be responsible for reducing SWA time [73]. Indeed,  $A\beta$

accumulation in the brain is accompanied by hyperactivity of the cortical neurons [73,74,75], and blocking neuron depolarization using gamma-aminobutyric acid A significantly improves NREM-SWA time in mice with AD [73]. Thus,  $A\beta$ -mediated synaptic abnormalities associated with neural hyperactivity can be another possible mechanism underlying SWA deficit in AD [73,74,75,76,77,78]. A better understanding of the astrocyte-mediated mechanisms of regulation of SWA time can open a new niche for novel pharmacological therapy in AD.

Thus, the BWRS is an important function of the sleeping brain that can be disturbed in animals and patients with different brain diseases, especially AD. Further studies of the mechanisms of sleep activation of the BWRS and injuries of these processes could offer a new understanding of the role of sleep in the etiology of neurological pathologies. These innovative findings could also contribute to the development of new technologies for the modulation of the therapeutic properties of the BWRS during deep sleep.

In general, significant progress in understanding the restorative mechanisms of sleep occurred in 2013, when the first study on the activation of the BWRS during deep sleep in mice was published [15]. The use of new optical technology, multiphoton microscopy, made it possible to study the movement of brain fluids in living mice with

simultaneous EEG recording.

Later, using multiphoton microscopy, the glymphatic system was discovered, which is believed to have the function of removing metabolites and toxins and to be activated during sleep [79,80]. However, due to intensive research in this area, it has become clear that the perivascular spaces are more preferable routes for the removal of unnecessary compounds, including  $A\beta$ , from the brain than the glymphatic pathway [81]. In recent human studies, it was confirmed that deep sleep is accompanied by an increase in CSF generation and the activation of the BWRS [23]. These pioneering results are an important informative platform for the development of fundamentally new technologies for studying and controlling the restorative mechanisms of sleep [9,28,82,83]. Pilot studies on rats show that sleep and opened BBB are equally accompanied by BWRS activation, which has an identical effect on the EEG dynamics and makes it possible to identify EEG markers of increased BBB permeability [84]. Non-invasive technology for assessing the BBB using only EEG is important for monitoring BBB permeability during surgery in order to control the depth of anesthesia and for the analysis of the progression of a number of brain diseases, such as AD and Parkinson's disease, brain trauma, diabetic and COVID-19 cerebrovascular injuries, and depression [16,85,86,87,88,89,90,91,92,93,94,95,96,97,98].

## **2. Photobiomodulation of BWRS: Innovative Strategies for Night Therapy of Brain Diseases**

The meningeal lymphatic vessels (MLVs) are regarded as an important part of the BWRS [99,100,101,102,103,104,105]. BWRS dysfunction due to MLV abnormalities contributes to the development of neurodegenerative diseases, brain tumors and cerebrovascular accidents [61,62,63,102,103,104,105,106,107,108,109,110]. Emerging evidence indicates that MLVs have attracted a lot of therapeutic interest. However, currently, there are no clinically approved non-pharmacological technologies for the modulation of MLV function. Recent studies suggest that transcranial photobiomodulation (tPBM) of MLVs might be an innovative technology to target the BWRS [9,28,82,83,111,112,113,114,115].

Recent reviews highlight the actual applications of tPBM to the therapy of sleep disorders [9,116] and brain diseases associated with sleep injuries, including AD [117,118,119,120], psychological problems [121], brain trauma [122] and other neurological diseases [123,124]. tPBM, known as low-level light therapy, is a technique of non-invasive treatment with red and near-infrared light irradiation in a photo-therapeutic window (between 600 and 1300 nm) [125,126]. However, light at short wavelengths has

very limited penetration into the brain due to light scattering, while light at long wavelengths has heating effects due to tissue absorption [126]. Light at wavelengths within range of 600–1300 nm has maximum penetration depth into the brain because of its minimal absorption and scattering within biological tissues; thus, it has significant PBM capability [125,126,127,128]. The light wavelength of 1300 nm has less scattering and can penetrate deeper into the brain [128]. Recent studies on mice show that tPBM (1267 nm) stimulates A $\beta$  removal from the brain, which contributes to the improvement of their neurological status [28,129]. Li et al. demonstrated the therapeutic effects of tPBM (1267 nm) in mice in a model of intraventricular hemorrhages (IVHs) [130]. The course of tPBM accelerated red blood cell evacuation from the ventricles, which improved the neurological outcome and reduced mortality in mice [130]. tPBM (1267 nm) could also be a promising technology for the modulation of the lymphatic delivery of drugs and nanocarriers to the brain pathology bypassing the BBB [115]. These pilot findings suggest that tPBM at the light wavelength of 1267 nm could be a novel, non-invasive, readily applicable and commercially viable technology for the routine treatment of various brain diseases.

tPBM of the brain has been studied extensively for prolonged periods and numerous devices are available on the market [123]. Figure 2a illustrates a typical tPBM

procedure, which includes irradiation of the patient's head with an expanded 1064 nm laser beam [131]. Laser light radiation is applied with a handheld laser head, controlled by a physician [131,132,133], because of the potential hazard of invisible collimated laser beams and the requirement to use appropriate eye and skin protection. Light sources based on light-emitting diodes (LEDs) provide more flexible applications of tPBM, as they are safer for patients and cheaper than a laser of equivalent power density. To ensure uniform light energy distribution over a skin surface, LEDs are assembled into clusters of various sizes and constructions, rigid blocks [118,134] or flexible panels [135]. LED clusters can be fixed on a patient's head with a headband [136], mounted inside a rigid helmet [134], or assembled in a helmet-like structure [137] or in a hat made of flexible LED panels [135] (Figure 2a–d).

**Figure 2.**

[Open in a new tab](#)

Devices for tPBM: **(a)** typical tPBM procedure includes irradiation of patient' head with expanded 1064 nm laser beam [131], which has maximum penetration depth into the brain [125,126,127,128]; **(b)** helmet for tPBM with system of forced cooling [134]; **(c)** helmet with

rigid structure assembled of 1070 nm LED clusters [137]; (d) hat made of flexible LED panels [135]; (e) smart sleep glasses [138].

To prevent the heating effect on the head, tPBM devices assembled with a large number of LEDs are used with a cooling system mounted over each LED cluster [134,137] (Figure 2b). Other therapeutic applications of PBM include the modification of ambient light with blue light emitted with an eyeglass-shaped gadget to correct delayed wake phase disorder [138] (Figure 2e).

Typically, the tPBM procedure is performed on awake patients. However, BWRS activity is much stronger during sleep than in the awake state, as we discuss above [15,28,29,30,31,32]. Based on the fact that the BWRS is activated during deep (NREM) sleep, we hypothesize that tPBM during night sleep might be more effective for the therapy of brain diseases than tPBM during the day. In our recent review, we discuss that sleep can be a promising therapeutic target for cerebral small-vessel disease [9]. In our pioneering study, we discovered that tPBM performed during sleep stimulates A $\beta$  clearance from the mouse brain more effectively than tPBM applied during wakefulness [28]. The course of night tPBM promotes good recovery of neurological status and recognition memory in mice with AD compared with the daily course of tPBM [28]. Several reviews report that the mechanisms of therapeutic effects of tPBM during sleep could be completely different from those

during the awake state [83,139]. However, the application of tPBM during sleep as a new strategy for the night therapy of brain diseases is in its infancy. Although the demand for tPBM during sleep has been clearly formulated, its practical realization has not yet been reported [83,139].

Currently, there are no technologies for simultaneous tPBM and sleep monitoring. Recent surveys of consumer sleep technologies demonstrate some demand for optional physical interventions during the sleep process aimed to improve the quality of sleep [140], although the requirement of strong scientific evidence to legitimize claims about utility, safety, and efficacy, as well as for informed choice and public trust, are emphasized [141,142]. Sleep technologies are presented on the market as various sleep-tracking devices [143,144] (Figure 3a–d).

**Figure 3.**

[Open in a new tab](#)

Sleep-tracking gadgets: **(a)** BrainBit headband [143]; **(b)** Philips Smart Sleep Deep Sleep Headband [142]; **(c)** Oura ring smart sleep tracker [144]; **(d)** sleep tracker 2 mm thick belt to be placed under the sheet on a bed [144].

Commercial sleep-tracking technologies include radars; near-infrared and thermal imaging devices; bedding-based sleep sensors; and wearable body sensors designed to be fixed on wrist, finger, feet, waist, chest or head [143,145]. Headbands are also capable of performing

electroencephalography (EEG) recording and analysis that sufficiently improve sleep detection [142,144]. All sleep-tracking technologies use wireless interfaces to communicate with mobile devices utilizing applications for sleep monitoring.

### **3. Mechanisms of tPBM of BWRS**

There are a number of studies on the mechanisms underlying tPBM of the BWRS [9,28,111,112,113,114,115,129]. It is reported that PBM causes the dilation of both MLVs and mesenteric lymphatics [114,130]. The PBM-mediated dilation of lymphatic vessels (LVs) is associated with an increase in the permeability of the lymphatic endothelium and a decrease in the expression of tight junction (TJ) proteins [114]. TJ proteins are the main components of the lymphatic endothelium and play a crucial role in the regulation of lymph movement in the lymphatic network [146]. Changes in the permeability of the lymphatic endothelium allow wastes and immune cells to be transported with lymph in LVs [147,148]. Indeed, the transport of antigens, and immune and dendritic cells across LVs is coupled with water flux and depends on the permeability of LVs [149,150]. These effects of PBM might be related to the PBM-induced production of nitric oxide (NO) in the lymphatic endothelium [82,130,151]. NO dilates blood and lymphatic vessels via the activation of soluble

guanylate cyclase and protein kinase G, which stimulates the opening of calcium ( $\text{Ca}^{2+}$ )-activated potassium channels [152]. The reduction in the  $\text{Ca}^{2+}$  intracellular level blocks the phosphorylation of myosin light-chain kinase, leading to vascular relaxation [152]. There are several studies suggesting that PBM-mediated vascular relaxation is the result of a PBM-induced increase in the activity of endothelial NO synthase (eNOS) [82,153,154,155]. eNOS-induced NO production in the lymphatic endothelium leads to an increase in lymphatic flow and the removal of wastes and toxins from different tissues [156].

Recently, it has been shown that PBM increases lymphatic contractility and that PBM stimulates NO synthesis in isolated cells of the lymphatic endothelium [115,130]. Lymphatic contractility is important for the movement of cells and molecules in LVs and is regulated by NO [157]. These facts suggest that the PBM-induced lymphatic contractility could be the possible mechanism responsible for the PBM-mediated increase in the clearance of wastes and metabolites from the brain [112,115,130].

Another mechanism of PBM-mediated control of vascular tone is the ability of singlet oxygen to regulate endothelial relaxation [158]. PBM at the light wavelength of 1267 nm stimulates the direct generation of singlet oxygen in tissues [159,160,161,162]. Singlet oxygen induces the oxidation of

the amino acid tryptophan in mammalian tissues, which leads to cell production of metabolites such as N-formylkynurenine with the activation of a heme-containing enzyme called indoleamine 2, 3-dioxygenase 1 [158]. This enzyme is widely expressed in the blood and lymphatic endothelium, contributing to the relaxation of vascular tone [158,163,164]. It was recently discovered that endothelial indoleamine 2, 3-dioxygenase 1 causes the dilation of blood vessels via the formation of singlet oxygen [158]. These pioneering findings shed light on the role of singlet oxygen in the regulation of vascular tone, opening a new perspective on the modulation of the systemic inflammatory response of the vasculature.

The PBM-mediated increase in NO production inhibits the activity of the mitochondrial cytochrome c oxidase (CCO) enzyme, leading to an increase in mitochondrial membrane potential [165,166,167]. These PBM effects contribute to an increase in the metabolic and energetic activity of cells via more oxygen consumption, more glucose metabolization and more ATP production by the mitochondria [165,166,167,168]. Recently, a mechanism of PBM-mediated stimulation of A $\beta$  clearance from the brain via the CCO-mediated activation of the cAMP-dependent protein kinase signaling pathway was discovered [169]. The neuroprotective effects of PBM (1068 nm) due to a decrease in the A $\beta$  level in the brain through microglia activation and

angiogenesis were also demonstrated in mice with AD [170]. PBM decreases A $\beta$  deposition in the brain by recruiting microglia to the A $\beta$  burden [170]. PBM can reduce A $\beta$  accumulation in the brain by increasing angiogenesis [171]. Indeed, the PBM (1070 nm)-induced increase in cerebral vessel density is positively correlated with the clearance of toxins from the brain [171]. PBM (630 nm) also improves brain drainage by contributing to A $\beta$  removal in the APP/PS1 mouse model of AD [172].

## **4. Conclusions**

In modern neurobiology, sleep is considered a novel biomarker and a promising therapeutic target for brain diseases. This is due to the recent discovery of the nighttime activation of the BRWS, which plays an important role in the removal of wastes and toxins from the brain, contributing to the neuroprotection of the CNS. The hypothesis has arisen in the scientific community that night stimulation of the BWRS might be a breakthrough strategy as a new treatment of AD, Parkinson's disease, stroke, brain trauma and oncology. Although this research is in its infancy, there are pioneering and promising results suggesting that night tPBM stimulates the lymphatic removal of A $\beta$  from the mouse brain more effectively than daily tPBM and is associated with a greater improvement of the neurological status and recognition memory in animals. There is the hypothesis that

tPBM modulates the tone and permeability of the lymphatic endothelium by stimulating NO formation, promoting the lymphatic clearance of wastes and toxins from brain tissues (Figure 4). It is assumed that tPBM can also lead to angio- and lymphangiogenesis, which might be another mechanism responsible for tPBM-mediated stimulation of the BWRS. Typically, tPBM is used in the awake state. Therefore, there are numerous tPBM devices constructed to apply a certain dose of light irradiation without technologies for sleep monitoring. To create a technology for simultaneous tPBM and sleep monitoring, only a wirelessly controlled gadget with LED or laser sources is needed to perform tPBM during sleep. The gadget can be controlled directly using a sleep-tracking device or mobile application that integrates data from various sources, including ambient conditions, e.g., room temperature, atmospheric pressure etc. Thus, there is a crucial challenge to design autonomous LED or laser light sources that are capable of providing the required therapeutic dose of light radiation in a certain region of the patient's head without disturbing the sleeping patient, e.g., sleep-tracking headbands [142,144]. To minimize patient discomfort, advanced materials such as flexible organic LEDs can be used [173,174].

**Figure 4.**



[Open in a new tab](#)

Emerging strategy of tPBM of the BRWS during deep sleep (NREM-SWA activity with delta rhythm): **(a)** Since the BRWS is activated during deep sleep, tPBM-mediated stimulation of the clearance of wastes and toxins from the brain could be more effective if tPBM were used during the NREM-SWA activity of the sleeping brain than during wakefulness [28]; **(b–d)** There is the hypothesis that tPBM modulates the tone and permeability of the lymphatic endothelium via the stimulation of eNOS and NO production, promoting the lymphatic clearance of wastes and toxins from brain tissues [96,137,138,139,140].

Since the idea of tPDT of the BWRS during sleep is in its infancy, here, we only discuss animal studies, because there are no human data in this area yet. There is evidence of a possible pharmacological stimulation of lymphaneogenesis for the treatment of brain tumors and for increasing brain immunity, as well as for the lymphatic excretion of A $\beta$  from brain tissues [175,176]. However, bringing new pharmaceuticals to the market takes 10 to 15 years. The proposed new pharmacological strategies for the stimulation of the BWRS have been tested on a small group of animals. It could take a long time for additional preclinical and clinical trials. In addition, pharmacological studies are not always successful, as in the case of the development of aducanumab, a novel anti-A $\beta$  antibody for the treatment of AD. Two phase 3 clinical trials including 3285 participants demonstrated that aducanumab causes vasogenic edema in 35% of patients associated with different symptoms, such as headache, confusion, dizziness and nausea, and that microhemorrhages occur in 19% of AD patients [176]. Therefore, the Food and Drug Administration (FDA) required

Biogen, the pharmacological company, to conduct an additional clinical trial to verify aducanumab's clinical benefits. If the trial fails to verify clinical benefits, the FDA may initiate proceedings to withdraw the approval of aducanumab.

The purpose of our review is to highlight non-pharmacological strategies for the stimulation of the BWRS that have a high potential to be introduced into routine clinical practice, as well as those that can become smart sleep gadgets used at home and while traveling without reducing the high pace of life of modern people. In this aspect, photonic technologies are the most promising, since they are already widely used in the clinical field for the treatment of various brain diseases [117,118,119,120,121,122,123,124]. tPBM of the BWRS is the only non-pharmacological technology proposed to activate the lymphatic excretion of metabolites and toxins from brain tissues [9,28,82,83,111,112,113,114,115,129]. Further studies of optimal doses and wavelengths of tPDT, and advantages and limitations of tPBM of the BWRS in patients of different ages and with various brain diseases could significantly help in the development of guidelines for its safe use in humans. The auditory effects can affect SWA time [177], which may also influence the BWRS, and could be an interesting subject of future research.

# **Author Contributions**

O.S.-G., J.K., D.Z. and T.P. initiated and supervised this review article; O.S.-G. wrote Abstract and Conclusions; O.S.-G., A.K., A.T., E.K. and E.S. prepared the section "Therapeutic properties of activation of the brain waste removal system (BWRS) during deep sleep"; D.Z., D.L. and T.Y. prepared the section "Photobiomodulation of the BWRS: the innovative strategies for the night therapy of brain diseases"; O.S.-G., I.F., I.B. and V.T. prepared the section "Mechanisms of tPBM of the BWRS and the MLVs"; all authors were also involved in the editing of the manuscript; E.K. and E.S. prepared the figures. All authors have read and agreed to the published version of the manuscript.

## **Institutional Review Board Statement**

Not applicable.

## **Informed Consent Statement**

Not applicable.

## **Data Availability Statement**

Not applicable.

## **Conflicts of Interest**

The authors declare that they have no competing interest.

## Funding Statement

O.S.-G., A.T., A.K., I.B., V.T., J.K., T.P., E.S. and E.K. were supported by RF Governmental grant No. 075-15-2022-1094, grant Nos. 20-15-00090 and 21-75-10088 from RSF and grant Nos. 20-015-00308-a and China-a 19-515-55016 from RFBR. D.Z. was supported by National Natural Science Foundation of China (NSFC) (grant Nos. 61860206009 and 81870934) Innovation Fund of WNLO and Innovation Project of Optics Valley Laboratory (grant No. OVL2021BG011). T.Y. was supported by National Natural Science Foundation of China (NSFC) (grant No. 81961138015). D.L. was supported by National Natural Science Foundation of China (NSFC) (grant No. 82001877).

## Footnotes

**Disclaimer/Publisher's Note:** The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.

## References

1. Franks N.P., Wisden W. *The inescapable drive to sleep: Overlapping mechanisms of sleep and sedation.* *Science.* 2021;374:556–559. doi: 10.1126/science.abi8372.normal [DOI] [PubMed] [Google Scholar]
2. Sabia S., Fayosse A., Dumurgier J., Van Hees V.T., Paquet C., Sommerlad A., Kivimäki M., Dugravot A., Singh-Manoux A. *Association of sleep duration in middle and old age with incidence of dementia.* *Nat. Commun.* 2021;12:2289. doi: 10.1038/s41467-021-22354-2.normal [DOI] [PMC free article] [PubMed] [Google Scholar]
3. Ooms S., Overeem S., Besse K., Rikkert M.O., Marcel Verbeek M., Claassen J.A.H.R. *Effect of 1 night of total sleep deprivation on cerebrospinal fluid  $\beta$ -amyloid 42 in healthy middle-aged men: A randomized clinical trial.* *JAMA Neurol.* 2014;71:971–977. doi: 10.1001/jamaneurol.2014.1173.normal [DOI] [PubMed] [Google Scholar]
4. Shokri-Kojoria E., Wanga G.-J., Wiersa C.E., Demirala S.B., Guoa M., Kima S.W., Lindgren E., Ramirez V., Zehraa A., Freeman C., et al.  *$\beta$ -Amyloid accumulation in the human brain after one night of sleep deprivation.* *Proc. Natl. Acad. Sci. USA.* 2018;115:4483–4488. doi: 10.1073/pnas.1721694115.normal [DOI] [PMC free article] [PubMed] [Google Scholar]
5. Lucey B.P., McCullough A., Landsness E.C., Toedebusch C.D., McLeland J.S., Zaza A.M., Fagan A.M., McCue L., Xiong C., Morris J.C., et al. *Reduced non-rapid eye movement sleep is associated with tau pathology in early Alzheimer's*

disease. *Sci. Transl. Med.* 2019;11:eaau6550. doi:

10.1126/scitranslmed.aau6550.normal [[DOI](#)] [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]

6.Lutsey P.L., Misialek J.R., Mosley T.H., Gottesman R.F., Punjabi N.M., Shahar E., MacLehose R., Ogilvie R.P., Knopman D., Alonso A. Sleep characteristics and risk of dementia and Alzheimer's disease: The atherosclerosis risk in communities study. *Alzheimer Dement.* 2018;14:157–166. doi: 10.1016/j.jalz.2017.06.2269.normal [[DOI](#)] [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]

7.Mendelsohn A.R., Larrick J.W. Sleep facilitates clearance of metabolites from the brain: Glymphatic function in aging and neurodegenerative diseases. *Rejuven. Res.* 2013;16:518–523. doi: 10.1089/rej.2013.1530.normal [[DOI](#)] [[PubMed](#)] [[Google Scholar](#)]

8.Carvalho D.Z., St Louis E.K., Knopman D.S., Boeve D.B.F., Lowe V.J., Roberts R.O., Mielke M.M., Przybelski S.A., Machulda M.M., Petersen R.C., et al. Association of excessive daytime sleepiness with longitudinal beta-amyloid accumulation in elderly persons without dementia. *JAMA Neurol.* 2018;75:672–680. doi: 10.1001/jamaneurol.2018.0049.normal [[DOI](#)] [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]

9.Semyachkina-Glushkovskaya O., Postnov D., Penzel T., Kurths J. Sleep as a novel biomarker and a promising therapeutic target for cerebral small vessel disease: A review focusing on Alzheimer's disease and the blood-brain barrier.

*Int. J. Mol. Sci.* 2020;21:6293. doi:

10.3390/ijms21176293.normal [[DOI](#)] [[PMC free article](#)]

[[PubMed](#)] [[Google Scholar](#)]

10.Izci-Balserak B., Zhu B., Wang H., Bronas U.G., Gooneratne N.S. Independent associations between sleep duration, gamma gap, and cognitive function among older adults: Results from the NHANES 2013–2014. *Geriatr. Nurs.* 2022;44:1–7. doi: 10.1016/j.gerinurse.2021.12.019.normal

[[DOI](#)] [[PubMed](#)] [[Google Scholar](#)]

11.Spira A.P., Gamaldo A.A., An Y., Wu M.N., Simonsick E.M., Bilgel M., Zhou Y., Wong D.F., Ferrucci L., Resnick S.M. Self-reported sleep and beta-amyloid deposition in community-dwelling older adults. *JAMA Neurol.* 2013;70:1537–1543.

doi: 10.1001/jamaneurol.2013.4258.normal [[DOI](#)] [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]

12.Minakawa E.N., Wada K., Nagai Y. Sleep disturbance as a potential modifiable risk factor for Alzheimer's disease. *Int. J. Mol. Sci.* 2019;20:803. doi:

10.3390/ijms20040803.normal [[DOI](#)] [[PMC free article](#)]

[[PubMed](#)] [[Google Scholar](#)]

13.Song D., Zhou J., Ma J., Chang J., Qiu Y., Zhuang Z., Xiao H., Zeng L. Sleep disturbance mediates the relationship between depressive symptoms and cognitive function in older adults with mild cognitive impairment. *Geriatr. Nurs.* 2021;42:1019–1023. doi:

10.1016/j.gerinurse.2021.06.004.normal [[DOI](#)] [[PubMed](#)]

[[Google Scholar](#)]

14. Wang Z., Heizhati M., Wang L., Li M., Yang Z., Lin M., Abudereyimu R., Hong J., Yang W., Yao L., et al. Poor sleep quality is negatively associated with low cognitive performance in general population independent of self-reported sleep disordered breathing. *BMC Public Health*. 2022;22:3. doi: 10.1186/s12889-021-12417-w.normal [[DOI](#)] [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]

15. Xie L., Kang H., Xu Q., Chen M.J., Liao Y., Thiyagarajan M., O'Donnell J., Christensen D.J., Nicholson C., Iliff J.J., et al. Sleep drives metabolite clearance from the adult brain. *Science*. 2013;342:373–377. doi: 10.1126/science.1241224.normal [[DOI](#)] [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]

16. Semyachkina-Glushkovskaya O., Mamedova A., Vinnik V., Klimova M., Saranceva E., Ageev V., Yu T., Zhu D., Penzel T., Kurths J. Brain Mechanisms of COVID-19-Sleep Disorders. *Int. J. Mol. Sci.* 2021;22:6917. doi: 10.3390/ijms22136917.normal [[DOI](#)] [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]

17. Hurtado-Alvarado G., Becerril-Villanueva E., Contis-Montes O.A., Domínguez-Salazar E., Salinas-Jazmín N., Pérez-Tapia S.M., Pavon L., Velázquez-Moctezuma J., Gómez-González B. The yin/yang of inflammatory status: Blood-brain barrier regulation during sleep. *Brain Behav. Immun.* 2018;69:154–166. doi: 10.1016/j.bbi.2017.11.009.normal [[DOI](#)] [[PubMed](#)] [[Google Scholar](#)]

18. Pan W., Kastin A.J. *The blood-brain barrier: Regulatory roles in wakefulness and sleep. Neuroscientist.* 2017;23:124–136. doi: 10.1177/1073858416639005. normal [DOI] [PubMed] [Google Scholar]
19. Cuddapah V.A., Zhang S.L., Sehgal A. *Regulation of the blood-brain barrier by circadian rhythms and sleep. Trends Neurosci.* 2019;42:500–510. doi: 10.1016/j.tins.2019.05.001. normal [DOI] [PMC free article] [PubMed] [Google Scholar]
20. Medina-Flores F., Hurtado-Alvarado G., Contis-Montes de Oca A., López-Cervantes S.P., Konigsberg M., Deli M.A., Gómez-González B. *Sleep loss disrupts pericyte-brain endothelial cell interactions impairing blood-brain barrier function. Brain Behav. Immun.* 2020;89:118–132. doi: 10.1016/j.bbi.2020.05.077. normal [DOI] [PubMed] [Google Scholar]
21. Bellesi M., de Vivo L., Chini M., Gilli F., Tononi G., Chiara Cirelli C. *Sleep Loss Promotes Astrocytic Phagocytosis and Microglial Activation in Mouse Cerebral Cortex. J. Neurosci.* 2017;37:5263–5273. doi: 10.1523/JNEUROSCI.3981-16.2017. normal [DOI] [PMC free article] [PubMed] [Google Scholar]
22. Papachristou C.S. *Aristotle's Theory of 'Sleep and Dreams' in the light of Modern and Contemporary Experimental Research. Electron. J. Philos.* 2014;17:1–47. doi: 10.18267/j.e-logos.374. normal [DOI] [Google Scholar]
23. Fultz N.E., Bonmassar G., Setsompop K., Stickgold R.A.,

Rosen B.R., Polimeni J.R., Lewis L.D. Coupled electrophysiological, hemodynamic, and cerebrospinal fluid oscillations in human sleep. *Science*. 2019;366:628–631. doi: 10.1126/science.aax5440.normal [[DOI](#)] [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]

24.Reddy O.C., Van der Werf Y.D. *The Sleeping Brain: Harnessing the Power of the Glymphatic System through Lifestyle Choices*. *Brain Sci*. 2020;10:868. doi: 10.3390/brainsci10110868.normal [[DOI](#)] [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]

25.Silva I., Silva J., Ferreira R., Trigo D. Glymphatic system, AQP4, and their implications in Alzheimer's disease. *Neurol Res Pract*. 2021;3:5. doi: 10.1186/s42466-021-00102-7.normal [[DOI](#)] [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]

26.Yi T., Gao P., Zhu T., Yin H., Jin S. Glymphatic System Dysfunction: A Novel Mediator of Sleep Disorders and Headaches. *Front. Neurol*. 2022;13:885020. doi: 10.3389/fneur.2022.885020.normal [[DOI](#)] [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]

27.Yan T., Qiu Y., Yu X., Yang L. Glymphatic Dysfunction: A Bridge Between Sleep Disturbance and Mood Disorders. *Front. Psychiatry*. 2021;12:658340. doi: 10.3389/fpsyt.2021.658340.normal [[DOI](#)] [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]

28.Semyachkina-Glushkovskaya O., Penzel T., Blokhina I., Khorovodov A., Fedosov I., Yu T., Karandin G., Evsukova A., Elovenko D., Adushkina V., et al. Night photostimulation of

clearance of beta-amyloid from mouse brain: New strategies in preventing Alzheimer's disease. *Cells*. 2021;10:3289. doi: 10.3390/cells10123289.normal [[DOI](#)] [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]

29.Nedergaard M., Goldman S.A. Glymphatic failure as a final common pathway to dementia. *Science*. 2020;370:50–56. doi: 10.1126/science.abb8739.normal [[DOI](#)] [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]

30.Hablitz L.M., Vinitzky H.S., Sun Q., Stæger F.F., Sigurdsson B., Mortensen K.N., Lilius T.O., Nedergaard M. Increased glymphatic influx is correlated with high EEG delta power and low heart rate in mice under anesthesia. *Sci. Adv.* 2019;5:eaav5447. doi: 10.1126/sciadv.aav5447.normal [[DOI](#)] [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]

31.Benveniste H., Heerdt P.M., Fontes M., Rothman D.L., Volkow N.D. Glymphatic system function in relation to anesthesia and sleep states. *Anesth. Analg.* 2019;128:747–758. doi: 10.1213/ANE.00000000000004069.normal [[DOI](#)] [[PubMed](#)] [[Google Scholar](#)]

32.Lee H., Xie L., Yu M., Kang H., Feng T., Deane R., Logan J., Nedergaard M., Benveniste H. The effect of body posture on brain glymphatic transport. *J. Neurosci.* 2015;35:11034–11044. doi: 10.1523/JNEUROSCI.1625-15.2015.normal [[DOI](#)] [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]

33.Borb A.A., Achermann P. Sleep homeostasis and models of sleep regulation. *J. Biol. Rhythm.* 1999;14:559–570. doi: 10.1177/074873099129000894.normal [[DOI](#)] [[PubMed](#)]

[\[Google Scholar\]](#)

34. Tononi G., Cirelli C. *Sleep and the price of plasticity: From synaptic and cellular homeostasis to memory consolidation and integration.* *Neuron.* 2014;81:12–34. doi:

10.1016/j.neuron.2013.12.025.normal [\[DOI\]](#) [\[PMC free article\]](#) [\[PubMed\]](#) [\[Google Scholar\]](#)

35. Diekelmann S., Born J. *The memory function of sleep.* *Nat. Rev. Neurosci.* 2010;11:114–126. doi:

10.1038/nrn2762.normal [\[DOI\]](#) [\[PubMed\]](#) [\[Google Scholar\]](#)

36. Hauner K.K., Howard J.D., Zelano C., Gottfried J.A. *Stimulus-specific enhancement of fear extinction during slow-wave sleep.* *Nat. Neurosci.* 2013;16:1553. doi:

10.1038/nn.3527.normal [\[DOI\]](#) [\[PMC free article\]](#) [\[PubMed\]](#) [\[Google Scholar\]](#)

37. Lee Y.F., Gerashchenko D., Timofeev I., Bacskai B.J., Kastanenka K.V. *Slow wave sleep as a promising intervention target for Alzheimer's disease.* *Front. Neurosci.*

2020;14:705. doi: 10.3389/fnins.2020.00705.normal [\[DOI\]](#)

[\[PMC free article\]](#) [\[PubMed\]](#) [\[Google Scholar\]](#)

38. Niwa Y., Kanda G.N., Yamada R.G., Shi S., Sunagawa G.A., Ukai-Tadenuma M., Fujishima H., Matsumoto N., Masumoto K.H., Nagano M., et al. *Muscarinic acetylcholine receptors Chrm1 and Chrm3 are essential for REM sleep.* *Cell Rep.*

2018;24:2231–2247. doi:

10.1016/j.celrep.2018.07.082.normal [\[DOI\]](#) [\[PubMed\]](#)

[\[Google Scholar\]](#)

39. Funato H., Miyoshi C., Fujiyama T., Kanda T., Sato M.,

Wang Z., Ma J., Nakane S., Tomita J., Ikkyu A., et al. *Forward-genetics analysis of sleep in randomly mutagenized mice*. *Nature*. 2016;539:378–383. doi:

10.1038/nature20142.normal [[DOI](#)] [[PMC free article](#)]

[[PubMed](#)] [[Google Scholar](#)]

40.Cai X., Qiao J., Kulkarni P., Harding I.C., Ebong E., Ferris C.F. *Imaging the effect of the circadian light-dark cycle on the glymphatic system in awake rats*. *Proc. Natl. Acad. Sci. USA*. 2020;117:668–676. doi:

10.1073/pnas.1914017117.normal [[DOI](#)] [[PMC free article](#)]

[[PubMed](#)] [[Google Scholar](#)]

41.Hablitz L.M., Plá V., Giannetto M., Vinitsky H.S., Stæger F.F., Metcalfe T. *Circadian control of brain glymphatic and lymphatic fluid flow*. *Nat. Commun*. 2020;11:4411. doi:

10.1038/s41467-020-18115-2.normal [[DOI](#)] [[PMC free article](#)]

[[PubMed](#)] [[Google Scholar](#)]

42.Cederroth C.R., Albrecht U., Bass J., Brown S.A., Dyhrfeld-Johnsen J., Gachon F. *Medicine in the fourth dimension*. *Cell Metab*. 2019;30:238–250. doi:

10.1016/j.cmet.2019.06.019.normal [[DOI](#)] [[PMC free article](#)]

[[PubMed](#)] [[Google Scholar](#)]

43.Ulv Larsen S.M., Landolt H.P., Berger W., Nedergaard M., Knudsen G.M., Holst S.C. *Haplotype of the astrocytic water channel AQP4 is associated with slow wave energy regulation in human NREM sleep*. *PLoS Biol*.

2020;18:3000623. doi:

10.1371/journal.pbio.3000623.normal [[DOI](#)] [[PMC free](#)]

[article](#)] [[PubMed](#)] [[Google Scholar](#)]

44. Brancaccio M., Patton A.P., Chesham J.E., Maywood E.S., Hastings M.H. Astrocytes control circadian timekeeping in the suprachiasmatic nucleus via glutamatergic signaling.

*Neuron*. 2017;93:1420–1435. doi:

10.1016/j.neuron.2017.02.030.normal [[DOI](#)] [[PMC free](#)

[article](#)] [[PubMed](#)] [[Google Scholar](#)]

45. Cirelli C., Tononi G. The why and how of sleep-dependent synaptic down-selection. *Semin. Cell. Dev. Biol.*

2022;125:91–100. doi: 10.1016/j.semcdb.2021.02.007.normal

[[DOI](#)] [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]

46. Rodriguez A.V., Funk C.M., Vyazovskiy V.V., Nir Y., Tononi G., Cirelli C. Why does sleep slow-wave activity increase after extended wake? Assessing the effects of increased cortical firing during wake and sleep. *J. Neurosci.*

2016;36:12436–12447. doi: 10.1523/JNEUROSCI.1614-

16.2016.normal [[DOI](#)] [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]

47. de Vivo L., Bellesi M., Marshall W., Bushong E.A., Ellisman M.H., Tononi G., Cirelli C. Ultrastructural evidence for synaptic scaling across the wake/sleep cycle. *Science*.

2017;355:507–510. doi: 10.1126/science.aah5982.normal

[[DOI](#)] [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]

48. Iliff J.J., Wang M., Zeppenfeld D.M., Venkataraman A., Plog B.A., Liao Y., Deane R., Nedergaard M. Cerebral arterial pulsation drives paravascular CSF-interstitial fluid exchange in the murine brain. *J. Neurosci.* 2013;33:18190–18199. doi:

10.1523/JNEUROSCI.1592-13.2013.normal [[DOI](#)] [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]

49. *Bedussi B., Naessens D.M.P., de Vos J., Olde Engberink R., Wilhelmus M.M.M., Richard E., ten Hove M., Van Bavel E., Bakker E.N.P. Enhanced interstitial fluid drainage in the hippocampus of spontaneously hypertensive rats. Sci. Rep. 2017;7:744. doi: 10.1038/s41598-017-00861-x.normal* [[DOI](#)] [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]

50. *Hubbard J., Gent T.C., Hoekstra M.M.B. Rapid fast-delta decay following prolonged wakefulness marks a phase of wake-inertia in NREM sleep. Nat. Commun. 2020;11:3130. doi: 10.1038/s41467-020-16915-0.normal* [[DOI](#)] [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]

51. *Hoekstra M., Emmenegger Y., Hubbard J., Franken P. Cold-inducible RNA-binding protein (CIRBP) adjusts clock-gene expression and REM-sleep recovery following sleep deprivation. eLife. 2019;8:43400. doi: 10.7554/eLife.43400.normal* [[DOI](#)] [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]

52. *Harding E.C., Yu X., Miao A., Andrews N., Ma Y., Ye Z., Lignos L., Miracca G., Ba W., Yustos R., et al. A neuronal hub binding sleep initiation and body cooling in response to a warm external stimulus. Curr. Biol. 2018;28:2263–2273. doi: 10.1016/j.cub.2018.05.054.normal* [[DOI](#)] [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]

53. *Jung C.M., Melanson E.L., Frydendall E.J., Perreault L., Eckel R.H., Wright K.P. Energy expenditure during sleep,*

sleep deprivation and sleep following sleep deprivation in adult humans. *J. Physiol.* 2011;589:235–244. doi: 10.1113/jphysiol.2010.197517.normal [[DOI](#)] [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]

54.McGinty D., Szymusiak R. Keeping cool: A hypothesis about the mechanisms and functions of slow-wave sleep. *Trends Neurosci.* 1990;13:480–487. doi: 10.1016/0166-2236(90)90081-K.normal [[DOI](#)] [[PubMed](#)] [[Google Scholar](#)]

55.Burhan H., Cherian I. Brain Cooling and Cleaning: A New Perspective in Cerebrospinal Fluid (CSF) Dynamics. In: Bozzetto Ambrosi P., Ahmad R., Abdullahi A., Agrawal A., editors. *Cerebrovascular Diseases*. IntechOpen; London, UK: 2020. normal [[Google Scholar](#)]

56.Sundaram S., Hughes R.L., Peterson E., Müller-Oehring E.M., Brontë-Stewart H.M., Poston K.L. Establishing a framework for neuropathological correlates and glymphatic system functioning in Parkinson's disease. *Neurosci. Biobehav. Rev.* 2019;103:305–315. doi: 10.1016/j.neubiorev.2019.05.016.normal [[DOI](#)] [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]

57.Pu T., Zou W., Feng W., Zhang Y., Wang L., Wang H. Persistent malfunction of glymphatic and meningeal lymphatic drainage in a mouse model of subarachnoid hemorrhage. *Exp. Neurobiol.* 2019;28:104–118. doi: 10.5607/en.2019.28.1.104.normal [[DOI](#)] [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]

58.Chen J., Wang L., Xu H. Meningeal lymphatics clear

*erythrocytes that arise from subarachnoid hemorrhage. Nat. Commun. 2020;11:3159. doi: 10.1038/s41467-020-16851-z.normal* [[DOI](#)] [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]

59.Liu X., Gao C., Yuan J. *Subdural haematomas drain into the extracranial lymphatic system through the meningeal lymphatic vessels. Acta Neuropathol. Commun. 2020;8:16. doi: 10.1186/s40478-020-0888-y.normal* [[DOI](#)] [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]

60.Bolte A.C., Dutta A.B., Hurt M.E. *Meningeal lymphatic dysfunction exacerbates traumatic brain injury pathogenesis. Nat. Commun. 2020;11:4524. doi: 10.1038/s41467-020-18113-4.normal* [[DOI](#)] [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]

61.Hu X., Deng Q., Ma L. *Meningeal lymphatic vessels regulate brain tumor drainage and immunity. Cell Res. 2020;30:229–243. doi: 10.1038/s41422-020-0287-8.normal* [[DOI](#)] [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]

62.Ma Q., Schlegel F., Bachmann S.B. *Lymphatic outflow of cerebrospinal fluid is reduced in glioma. Sci. Rep. 2019;9:14815. doi: 10.1038/s41598-019-51373-9.normal* [[DOI](#)] [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]

63.Da Mesquita S., Louveau A., Vaccari A. *Functional aspects of meningeal lymphatics in ageing and Alzheimer's disease. Nature. 2018;560:185–191. doi: 10.1038/s41586-018-0368-8.normal* [[DOI](#)] [[PMC free article](#)] [[PubMed](#)]

[\[Google Scholar\]](#)

64.Kastanenka K.V., Calvo-Rodriguez M., Hou S.S., Zhou H., Takeda S., Arbel-Ornath M., Lariviere A., Lee Y.F., Kim A., Hawkes J.M., et al. Frequency-dependent exacerbation of Alzheimer's disease neuropathophysiology. *Sci. Rep.* 2019;9:8964. doi: 10.1038/s41598-019-44964-z.normal

[\[DOI\]](#) [\[PMC free article\]](#) [\[PubMed\]](#) [\[Google Scholar\]](#)

65.Kent B.A., Strittmatter S.M., Nygaard H.B. Sleep and EEG power spectral analysis in three transgenic mouse models of Alzheimer's disease: APP/PS1, 3xTgAD, and Tg2576. *J. Alzheimer Dis.* 2018;64:1325–1336. doi: 10.3233/JAD-180260.normal [\[DOI\]](#) [\[PMC free article\]](#) [\[PubMed\]](#) [\[Google Scholar\]](#)

66.Castano-Prat P., Perez-Mendez L., Perez-Zabalza M., Sanfeliu C., Giménez-Llort L., Sanchez-Vives M.V. Altered slow (<1 Hz) and fast (beta and gamma) neocortical oscillations in the 3xTg-AD mouse model of Alzheimer's disease under anesthesia. *Neurobiol. Ageing.* 2019;79:142–151. doi: 10.1016/j.neurobiolaging.2019.02.009.normal [\[DOI\]](#) [\[PubMed\]](#) [\[Google Scholar\]](#)

67.Mander B.A., Rao V., Lu B., Saletin J.M., Lindquist J.R., Ancoli-Israel S., Jagust W., Walker M.P. Prefrontal atrophy, disrupted NREM slow waves and impaired hippocampal-dependent memory in aging. *Nat. Neurosci.* 2013;16:357. doi: 10.1038/nn.3324.normal [\[DOI\]](#) [\[PMC free article\]](#) [\[PubMed\]](#) [\[Google Scholar\]](#)

68.Mander B.A., Marks S.M., Vogel J.W., Rao V., Lu B.,

Saletin J.M., Ancoli-Israel S., Jagust W.J., Walker M.P. *b-amyloid disrupts human NREM slow waves and related hippocampus-dependent memory consolidation*. *Nat. Neurosci.* 2015;18:1051–1057. doi: 10.1038/nn.4035.normal [DOI] [PMC free article] [PubMed] [Google Scholar]

69. Westerberg C.E., Mander B.A., Florczak S.M., Weintraub S., Mesulam M.M., Zee P.C., Paller K.A. *Concurrent impairments in sleep and memory in amnesic mild cognitive impairment*. *J. Int. Neuropsychol. Soc. JINS.* 2012;18:490. doi: 10.1017/S135561771200001X.normal [DOI] [PMC free article] [PubMed] [Google Scholar]

70. Winer J.R., Mander B.A., Helfrich R.F., Maass A., Harrison T.M., Baker S.L., Knight R.T., Jagust W.J., Walker M.P. *Sleep as a potential biomarker of tau and b-amyloid burden in the human brain*. *J. Neurosci.* 2019;39:6315–6324. doi: 10.1523/JNEUROSCI.0503-19.2019.normal [DOI] [PMC free article] [PubMed] [Google Scholar]

71. O'Donnell J., Zeppenfeld D., McConnell E., Pena S., Nedergaard M. *Norepinephrine: A neuromodulator that boosts the function of multiple cell types to optimize CNS performance*. *Neurochem. Res.* 2012;37:2496–2512. doi: 10.1007/s11064-012-0818-x.normal [DOI] [PMC free article] [PubMed] [Google Scholar]

72. Cipolla M.J., Li R., Vitullo L. *Perivascular innervation of penetrating brain parenchymal arterioles*. *J. Cardiovasc. Pharmacol.* 2004;44:1–8. doi: 10.1097/00005344-200407000-00001.normal [DOI] [PubMed] [Google]

## [Scholar](#)]

73.Kastanenka K.V., Hou S.S., Shakerdge N., Logan R., Feng D., Wegmann S., Chopra V., Hawkes J.M., Chen X., Bacskai B.J. *Optogenetic restoration of disrupted slow oscillations halts amyloid deposition and restores calcium homeostasis in an animal model of Alzheimer's disease. PLoS ONE. 2017;12:e0170275. doi:*

*10.1371/journal.pone.0170275.normal* [[DOI](#)] [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]

74.Selkoe D.J. *Early network dysfunction in Alzheimer's disease. Science. 2019;365:540–541. doi:*

*10.1126/science.aay5188.normal* [[DOI](#)] [[PubMed](#)] [[Google Scholar](#)]

75.Cirrito J.R., Kang J.E., Lee J., Stewart F.R., Verges D.K., Silverio L.M., Bu G., Mennerick S., Holtzman D.M. *Endocytosis is required for synaptic activity-dependent release of amyloid- $\beta$  in vivo. Neuron. 2008;58:42–51. doi:*

*10.1016/j.neuron.2008.02.003.normal* [[DOI](#)] [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]

76.Kuchibhotla K.V., Lattarulo C.R., Hyman B.T., Bacskai B.J. *Synchronous hyperactivity and intercellular calcium waves in astrocytes in Alzheimer mice. Science. 2009;323:1211–1215. doi:*

*10.1126/science.1169096.normal* [[DOI](#)] [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]

77.Robinson S.R. *Neuronal expression of glutamine synthetase in Alzheimer's disease indicates a profound impairment of metabolic interactions with astrocytes.*

- Neurochem. Int. 2000;36:471–482. doi: 10.1016/S0197-0186(99)00150-3.normal [[DOI](#)] [[PubMed](#)] [[Google Scholar](#)]
- 78.Poskanzer K.E., Yuste R. Astrocytes regulate cortical state switching in vivo. *Proc. Natl. Acad. Sci. USA*. 2016;113:2675–2684. doi: 10.1073/pnas.1520759113.normal [[DOI](#)] [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]
- 79.Iliff J., Wang M., Liao Y., Plogg B., Peng W., Gundersen G., Benveniste H., Vates G., Deane R., Goldman S., et al. A paravascular pathway facilitates CSF flow through the brain parenchyma and the clearance of interstitial solutes, including amyloid  $\beta$ . *Sci. Transl. Med.* 2012;147:ra111. doi: 10.1126/scitranslmed.3003748.normal [[DOI](#)] [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]
- 80.Jessen N., Munk A., Lundgaard I., Nedergaard M. The Glymphatic System: A Beginner's Guide. *Neurochem. Res.* 2015;12:2583–2599. doi: 10.1007/s11064-015-1581-6.normal [[DOI](#)] [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]
- 81.Hladky S., Barrand M. The glymphatic hypothesis: The theory and the evidence. *Fluids Barriers CNS*. 2022;19:9. doi: 10.1186/s12987-021-00282-z.normal [[DOI](#)] [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]
- 82.Salehpour F., Khademi M., Bragin D., DiDuro J. Photobiomodulation therapy and the glymphatic system: Promising applications for augmenting the brain lymphatic drainage system. *Int. J. Mol. Sci.* 2022;23:2975. doi: 10.3390/ijms23062975.normal [[DOI](#)] [[PMC free article](#)]

[\[PubMed\]](#) [\[Google Scholar\]](#)

83.Cecile M., Valverde A., Dole M., Hoh Kam J. *The effect of photobiomodulation on the brain during wakefulness and sleep.* *Front. Neurosci.* 2022;16:1232. doi:

10.3389/fnins.2022.942536.normal [\[DOI\]](#) [\[PMC free article\]](#)

[\[PubMed\]](#) [\[Google Scholar\]](#)

84.Semyachkina-Glushkovskaya O., Karavaev A., Prokhorov M., Runnova A., Borovkova E., Ishbulatov Y., Hramkov A., Kulminskiy D., Semenova N., Sergeev K., et al. *EEG biomarkers of activation of the lymphatic drainage system of the brain during sleep and opening of the blood-brain barrier.* *Comput. Struct. Biotechnol. J.* 2023;21:758–768.

doi: 10.1016/j.csbj.2022.12.019.normal [\[DOI\]](#) [\[PMC free](#)

[article\]](#) [\[PubMed\]](#) [\[Google Scholar\]](#)

85.Spieth L., Berghoff S., Stumpf S., Winchenbach J., Michaelis T., Watanabe T., Gerndt N., Düking T., Hofer S., Ruhwedel T., et al. *Anesthesia triggers drug delivery to experimental glioma in mice by hijacking caveolar transport.* *Neurooncol. Adv.* 2021;3:vdab140. doi:

10.1093/noajnl/vdab140.normal [\[DOI\]](#) [\[PMC free article\]](#)

[\[PubMed\]](#) [\[Google Scholar\]](#)

86.Tétrault S., Chever O., Sik A., Amzica F. *Opening of the blood-brain barrier during isoflurane anaesthesia.* *Eur. J. Neurosci.* 2008;28:1330–1341. doi: 10.1111/j.1460-

9568.2008.06443.x.normal [\[DOI\]](#) [\[PubMed\]](#) [\[Google](#)

[Scholar\]](#)

87.Yang X., Chen X. *The Crosstalk between the Blood–Brain*

*Barrier Dysfunction and Neuroinflammation after General Anaesthesia. Curr. Issues Mol. Biol. 2022;44:5700–5717. doi: 10.3390/cimb44110386.normal [DOI] [PMC free article] [PubMed] [Google Scholar]*

88.Sweeney M., Sagare A., Zlokovic B. *Blood–brain barrier breakdown in Alzheimer disease and other neurodegenerative disorders. Nat. Rev. Neurol. 2018;14:133–150. doi: 10.1038/nrneurol.2017.188.normal [DOI] [PMC free article] [PubMed] [Google Scholar]*

89.Wang D., Chen F., Han Z., Yin Z., Ge X., Lei P. *Relationship between Amyloid- Deposition and Blood–Brain Barrier Dysfunction in Alzheimer’s Disease. Front. Cell. Neurosci. 2021;15:695479. doi: 10.3389/fncel.2021.695479.normal [DOI] [PMC free article] [PubMed] [Google Scholar]*

90.Al-Bachari S., Naish J., Parker G., Emsley H., Parkes L. *Blood–Brain Barrier Leakage Is Increased in Parkinson’s Disease. Front. Physiol. 2020;11:593026. doi: 10.3389/fphys.2020.593026.normal [DOI] [PMC free article] [PubMed] [Google Scholar]*

91.Pineda-Pardo J., Gasca-Salas C., Fernández-Rodríguez B., Rodríguez-Rojas R., Del Álamo M., Obeso I., Hernández-Fernández F., Trompeta C., Martínez-Fernández R., Matarazzo M., et al. *Striatal Blood-Brain Barrier Opening in Parkinson’s Disease Dementia: A Pilot Exploratory Study. Mov. Disord. 2022;37:2057–2065. doi: 10.1002/mds.29134.normal [DOI] [PubMed] [Google*

[Scholar](#)]

92. Wu Y., Wu H., Guo X., Pluimer B., Zhao Z. *Blood–Brain Barrier Dysfunction in Mild Traumatic Brain Injury: Evidence from Preclinical Murine Models.* *Front. Physiol.* 2020;11:1030. doi: 10.3389/fphys.2020.01030.normal

[[DOI](#)]

[[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]

93. Dion-Albert L., Cadoret A., Doney E., Kaufmann F.N., Dudek K.A., Daigle B., Parise L.F., Cathomas F., Samba N., Hudson N., et al. *Vascular and blood-brain barrier-related changes underlie stress responses and resilience in female mice and depression in human tissue.* *Nat. Commun.* 2022;13:164. doi: 10.1038/s41467-021-27604-x.normal

[[DOI](#)]

[[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]

94. Wu S., Yin Y., Du L. *Blood-Brain Barrier Dysfunction in the Pathogenesis of Major Depressive Disorder.* *Cell. Mol. Neurobiol.* 2022;42:2571–2591. doi: 10.1007/s10571-021-01153-9.normal

[[DOI](#)] [[PMC free article](#)] [[PubMed](#)] [[Google](#)

[Scholar](#)]

95. Prasad S., Sajja R., Naik P., Cucullo L. *Diabetes Mellitus and Blood-Brain Barrier Dysfunction: An Overview.* *J. Pharmacovigil.* 2014;2:125. doi: 10.4172/2329-

6887.1000125.normal

[[DOI](#)] [[PMC free article](#)] [[PubMed](#)]

[[Google Scholar](#)]

96. Banks W.A. *The Blood-Brain Barrier Interface in Diabetes Mellitus: Dysfunctions, Mechanisms and Approaches to Treatment.* *Curr. Pharm. Des.* 2020;26:1438–1447. doi:

10.2174/1381612826666200325110014.normal [[DOI](#)]

[\[PubMed\]](#) [\[Google Scholar\]](#)

97.Krasemann S., Haferkamp U., Pfefferle S., Woo M., Heinrich F., Schweizer M., Appelt-Menzel A., Cubukova A., Barenberg J., Leu J., et al. The blood-brain barrier is dysregulated in COVID-19 and serves as a CNS entry route for SARS-CoV-2. *Stem Cell Rep.* 2022;17:307–320. doi: 10.1016/j.stemcr.2021.12.011.normal [\[DOI\]](#) [\[PMC free article\]](#)

[\[PubMed\]](#) [\[Google Scholar\]](#)

98.Chen Y., Yang W., Chen F., Cui L. COVID-19 and cognitive impairment: Neuroinvasive and blood-brain barrier dysfunction. *J. Neuroinflamm.* 2022;19:222. doi: 10.1186/s12974-022-02579-8.normal [\[DOI\]](#) [\[PMC free article\]](#)

[\[PubMed\]](#) [\[Google Scholar\]](#)

99.Louveau A., Plog B.A., Antila S., Alitalo K., Nedergaard M., Kipnis J. Understanding the functions and relationships of the glymphatic system and meningeal lymphatics. *J. Clin. Investig.* 2017;127:3210–3219. doi: 10.1172/JCI90603.normal

[\[DOI\]](#) [\[PMC free article\]](#) [\[PubMed\]](#) [\[Google Scholar\]](#)

100.Da Mesquita S., Fu Z., Kipnis J. The meningeal lymphatic system: A new player in neurophysiology. *Neuron.* 2018;100:375–388. doi:

10.1016/j.neuron.2018.09.022.normal

[\[DOI\]](#) [\[PMC free article\]](#) [\[PubMed\]](#) [\[Google Scholar\]](#)

101.Semyachkina-Glushkovskaya O., Postnov D., Kurths J. Blood–brain barrier, lymphatic clearance, and recovery: Ariadne’s thread in labyrinths of hypotheses. *Int. J. Mol. Sci.* 2018;19:3818. doi: 10.3390/ijms19123818.normal [\[DOI\]](#)

[\[PMC free article\]](#) [\[PubMed\]](#) [\[Google Scholar\]](#)

102. Tavares G.A., Louveau A. Meningeal lymphatics: An immune gateway for the central nervous system. *Cells*. 2021;10:3385. doi: 10.3390/cells10123385.normal [\[DOI\]](#)

[\[PMC free article\]](#) [\[PubMed\]](#) [\[Google Scholar\]](#)

103. Louveau A., Herz J., Alme M.N., Salvador A.F., Dong M.Q., Viar K.E., Herod S.G., Knopp J., Setliff J.C., Lupi A.L., et al. CNS lymphatic drainage and neuroinflammation are regulated by meningeal lymphatic vasculature. *Nat. Neurosci.* 2018;21:1380–1391. doi: 10.1038/s41593-018-0227-9.normal [\[DOI\]](#) [\[PMC free article\]](#) [\[PubMed\]](#) [\[Google Scholar\]](#)

*Neurosci.* 2018;21:1380–1391. doi: 10.1038/s41593-018-0227-9.normal [\[DOI\]](#) [\[PMC free article\]](#) [\[PubMed\]](#) [\[Google Scholar\]](#)

[\[Google Scholar\]](#)

104. Ding X.B., Wang X.X., Xia D.H., Liu H., Tian H.Y., Fu Y., Chen Y.K., Qin C., Wang J.Q., Xiang Z., et al. Impaired meningeal lymphatic drainage in patients with idiopathic Parkinson's disease. *Nat. Med.* 2021;27:411–418. doi: 10.1038/s41591-020-01198-1.normal [\[DOI\]](#) [\[PubMed\]](#)

10.1038/s41591-020-01198-1.normal [\[DOI\]](#) [\[PubMed\]](#)

[\[Google Scholar\]](#)

105. Olate-Briones A., Escalona E., Salazar C., Herrada M.J., Liu C., Herrada A.A., Escobedo N. The meningeal lymphatic vasculature in neuroinflammation. *FASEB J.* 2022;36:e22276. doi: 10.1096/fj.202101574RR.normal [\[DOI\]](#)

2022;36:e22276. doi: 10.1096/fj.202101574RR.normal [\[DOI\]](#)

[\[PubMed\]](#) [\[Google Scholar\]](#)

106. Song E., Mao T., Dong H., Boisserand L.S.B., Antila S., Bosenberg M., Alitalo K., Thomas J.L., Iwasaki A. VEGF-C-driven lymphatic drainage enables immunosurveillance of brain tumours. *Nature*. 2020;577:689–694. doi:

10.1038/s41586-019-1912-x.normal [[DOI](#)] [[PMC free article](#)]

[[PubMed](#)] [[Google Scholar](#)]

107.Lan Y.L., Wang H., Chen A., Zhang J. Update on the current knowledge of lymphatic drainage system and its emerging roles in glioma management. *Immunology*.

2022;10:111. doi: 10.1111/imm.13517.normal [[DOI](#)] [[PubMed](#)]

[[Google Scholar](#)]

108.Yanev P., Poinsatte K., Hominick D., Khurana N., Zurbier K.R., Berndt M., Plautz E.J., Dellinger M.T., Stowe A.M. Impaired meningeal lymphatic vessel development worsens stroke outcome. *J. Cereb. Blood Flow Metab.*

2020;40:263–275. doi: 10.1177/0271678X18822921.normal

[[DOI](#)] [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]

109.Harrison I.F., Ismail O., Machhada A., Colgan N., Ohene Y., Nahavandi P., Ahmed Z., Fisher A., Meftah S., Murray

T.K., et al. Impaired glymphatic function and clearance of tau in an Alzheimer's disease model. *Brain*. 2020;143:2576–

2593. doi: 10.1093/brain/awaa179.normal [[DOI](#)] [[PMC free](#)

[article](#)] [[PubMed](#)] [[Google Scholar](#)]

110.Lemprière S. Meningeal lymphatic flow slows after mild traumatic brain injury. *Nat. Rev. Neurol.* 2020;16:600. doi:

10.1038/s41582-020-00420-5.normal [[DOI](#)] [[PubMed](#)]

[[Google Scholar](#)]

111.Zhinchenko E., Klimova M., Mamedova A., Agranovich I., Blokhina I., Antonova T., Terskov A., Shirokov A., Navolkin N., Morgun A., et al. Photostimulation of extravasation of beta-amyloid through the model of blood-brain barrier.

*Electronics*. 2020;9:1056. doi:

10.3390/electronics9061056.normal [[DOI](#)] [[Google Scholar](#)]

112.Semyachkina-Glushkovskaya O., Postnov D., Lavrova A., Fedosov I., Borisova E., Nikolenko V., Penzel T., Kurths J., Tuchin V. Biophotonic strategies of measurement and stimulation of the cranial and the extracranial lymphatic drainage function. *IEEE J. Sel. Top. Quantum Electron*. 2021;27:7400313. doi:

10.1109/JSTQE.2020.3045834.normal [[DOI](#)] [[Google Scholar](#)]

113.Semyachkina-glushkovskaya O., Abdurashitov A., Dubrovsky A. Photobiomodulation of lymphatic drainage and clearance: Perspective strategy for augmentation of meningeal lymphatic functions. *BOE*. 2020;11:725–734. doi:

10.1364/BOE.383390.normal [[DOI](#)] [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]

114.Semyachkina-Glushkovskaya O., Abdurashitov A., Klimova M. Photostimulation of cerebral and peripheral lymphatic functions. *Transl. Biophoton*.

2020;2:e201900036. doi: 10.1002/tbio.201900036.normal [[DOI](#)] [[Google Scholar](#)]

115.Semyachkina-Glushkovskaya O., Fedosov I., Shirokov A. Photomodulation of lymphatic delivery of liposomes to the brain bypassing the blood-brain barrier: New perspectives for glioma therapy. *Nanophotonics*. 2021;10:3215–3227. doi:

10.1515/nanoph-2021-0212.normal [[DOI](#)] [[Google Scholar](#)]

116.Abad V., Guillemineault C. Diagnosis and treatment of

*sleep disorders: A brief review for clinicians. Dialog Clin.*

*Neurosci. 2022;5:371–388. doi:*

*10.31887/DCNS.2003.5.4/vabad.normal* [[DOI](#)] [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]

117. *Wu C., Yang L., Feng S., Zhu L. Therapeutic non-invasive brain treatments in Alzheimer's disease: Recent advances and challenges. Inflamm. Regen. 2022;42:31. doi:*

*10.1186/s41232-022-00216-8.normal* [[DOI](#)] [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]

118. *Hamblin M.R. Photobiomodulation for Alzheimer's disease: Has the light dawned? Photonics. 2019;6:77. doi:*

*10.3390/photonics6030077.normal* [[DOI](#)] [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]

119. *Cardoso F., Gonzalez-Lima F., Gomes da Silva S.*

*Photobiomodulation for the aging brain. Ageing Res. Rev.*

*2021;70:101415. doi: 10.1016/j.arr.2021.101415.normal* [[DOI](#)] [[PubMed](#)] [[Google Scholar](#)]

120. *Liu Y.-L., Gong S.Y., Shu-Ying X., Wang Y., Peng H.,*

*Shen Y., Liu C.-F. Light therapy: A new option for*

*neurodegenerative diseases. Chin. Med. J. 2021;134:634–*

*645. doi: 10.1097/CM9.0000000000001301.normal* [[DOI](#)] [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]

121. *Gutiérrez-Menéndez A., Marcos-Nistal M., Mendez M.,*

*Arias J. Photobiomodulation as a promising new tool in the management of psychological disorders: A systematic review. Neurosci. Biobehav. Rev. 2020;119:242–254. doi:*

*10.1016/j.neubiorev.2020.10.002.normal* [[DOI](#)] [[PubMed](#)]

[\[Google Scholar\]](#)

122.Srisurapanont K., Samakarn Y., Kamklong B., Siratrairat P. *Blue-wavelength light therapy for post-traumatic brain injury sleepiness, sleep disturbance, depression, and fatigue: A systematic review and network meta-analysis.* *PLoS ONE.* 2021;16:e0246172. doi:

10.1371/journal.pone.0246172.normal [\[DOI\]](#) [\[PMC free article\]](#) [\[PubMed\]](#) [\[Google Scholar\]](#)

123.Hamblin M.R., Huang Y.Y. *Photobiomodulation in the Brain: Low-Level Laser (Light) Therapy in Neurology and Neuroscience.* Academic Press; London, UK: 2019. 26p normal [\[Google Scholar\]](#)

124.Mosilhy E.A., Alshial E., Eltaras M.M., Rahman M.M.A., Helmy H.I., Elazoul A.H., Hamdy O., Haitham S.M. *Non-invasive transcranial brain modulation for neurological disorders treatment: A narrative review.* *Life Sci.*

2022;307:120869. doi: 10.1016/j.lfs.2022.120869.normal [\[DOI\]](#) [\[PubMed\]](#) [\[Google Scholar\]](#)

125.Kim M.M., Darafsheh A. *Light Sources and Dosimetry Techniques for Photodynamic Therapy.* *Photochem. Photobiol.* 2020;96:280–294. doi: 10.1111/php.13219.normal [\[DOI\]](#) [\[PubMed\]](#) [\[Google Scholar\]](#)

126.Tedford C.E., DeLapp S., Jacques S., Anders J. *Quantitative analysis of transcranial and intraparenchymal light penetration in human cadaver brain tissue.* *Lasers Surg. Med.* 2015;47:312–322. doi: 10.1002/lsm.22343.normal [\[DOI\]](#) [\[PubMed\]](#) [\[Google Scholar\]](#)

127. Salehpour F., Cassano P., Rouhi N., Hamblin M.R., De Taboada L., Farajdokht F., Mahmoudi J. Penetration profiles of visible and near-infrared lasers and light-emitting diode light through the head tissues in animal and human species: A review of literature. *Photobiomodul. Photomed. Laser Surg.* 2019;37:581–595. doi:

10.1089/photob.2019.4676.normal [[DOI](#)] [[PubMed](#)] [[Google Scholar](#)]

128. Wang T., Ouzounov D., Wu C., Horton N.G., Zhang B., Wu C.-H. Three-photon imaging of mouse brain structure and function through the intact skull. *Nat. Methods.* 2018;15:789–792. doi: 10.1038/s41592-018-0115-y.normal

[[DOI](#)] [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]

129. Zinchenko E., Navolokin N., Shirokov A., Khlebtsov B., Dubrovsky A., Saranceva E., Abdurashitov A., Khorovodov A., Terskov A., Mamedova A. Pilot study of transcranial photobiomodulation of lymphatic clearance of beta-amyloid from the mouse brain: Breakthrough strategies for non-pharmacologic therapy of Alzheimer's disease. *BOE.* 2019;10:4003–4017. doi: 10.1364/BOE.10.004003.normal

[[DOI](#)] [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]

130. Li D., Liu S., Yu T., Liu Z., Sun S., Bragin D., Navolokin N., Kurths J., Glushkovskaya-Semyachkina O., Zhu D.

Photostimulation of lymphatic clearance of red blood cells from the mouse brain after intraventricular hemorrhage.

*bioRxiv.* 2020;16:384149.normal [[Google Scholar](#)]

131. Vargas E., Barratt D.W., Saucedo C.L., Huang L.-D.,

Abraham J.A., Tanaka H., Haley A.P., Gonzales-Lima F. Beneficial neurocognitive effects of transcranial laser in older adults. *Lasers Med. Sci.* 2017;32:1153–1162. doi: 10.1007/s10103-017-2221-y.normal [[DOI](#)] [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]

132.Wang X., Wanniarachchi H., Wu A., Gonzalez-Lima F. Transcranial photobiomodulation and thermal stimulation induce distinct topographies of EEG alpha and beta power changes in healthy humans. *Sci. Rep.* 2021;11:18917. doi: 10.1038/s41598-021-97987-w.normal [[DOI](#)] [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]

133.Shahdadian S., Wang X., Wanniarachchi H., Chaudhari A., Truong N., Liu H. Neuromodulation of brain topography and network topology by prefrontal transcranial photobiomodulation. *J. Neural Eng.* 2022;19:066013. doi: 10.1088/1741-2552/ac9ede.normal [[DOI](#)] [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]

134.Longo M.G.F., Can Ozan Tan C.O., Chan S.-T., Welt J. Effect of transcranial low-level light therapy vs sham therapy among patients with moderate traumatic brain injury: A randomized clinical trial. *JAMA Netw. Open.* 2020;3:e2017337. doi: 10.1001/jamanetworkopen.2020.17337.normal [[DOI](#)] [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]

135.Hipskind S.G., Grover F.L., Hipskind S.G., Fort T.R. Pulsed transcranial red/near-infrared light therapy using light-emitting diodes improves cerebral blood flow and

cognitive function in veterans with chronic traumatic brain injury: A case series. *Photomed. Laser Surg.* 2018;37:77–84. doi: 10.1089/photob.2018.4489.normal [[DOI](#)] [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]

136.Cassano P., Mischoulon D., Cusin C., Petrie S. Transcranial photobiomodulation for the treatment of major depressive disorder. The ELATED-2 pilot trial. *Photomed. Laser Surg.* 2018;36:634–646. doi: 10.1089/pho.2018.4490.normal [[DOI](#)] [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]

137.Berman M.H., Halper J., Trent N., Jarrett H. Photobiomodulation with near infrared light helmet in a pilot, placebo controlled clinical trial in dementia patients testing memory and cognition. *J. Neurol. Neurosci.* 2017;8:176. doi: 10.21767/2171-6625.1000176.normal [[DOI](#)] [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]

138.Dong L., Fang P., Liu H., Chen L., Fu Y., Xie B., Liu Y., Ye H., Gu P. The clinical effect of blue light therapy on patients with delayed sleep-wake phase disorder. *Nat. Sci. Sleep.* 2022;14:75. doi: 10.2147/NSS.S344616.normal [[DOI](#)] [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]

139.Valverde A., Mitrofanis J., Moro C., Hamilton C. Lights at night: Does photobiomodulation improve sleep? *Neural Regen. Res.* 2022;18:474–477. doi: 10.4103/1673-5374.350191.normal [[DOI](#)] [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]

140.Devine J.K., Schwartz L.P., Choynowski J., Hursh S.R.

- Expert Demand for Consumer Sleep Technology Features and Wearable Devices. IoT. 2021;2:18. doi: 10.3390/iot3020018.normal* [[DOI](#)] [[Google Scholar](#)]
141. *McCall I.C., Lua C., Minielly N., Illis J. Owning ethical innovation: Claims about commercial wearable brain technologies. Neuron. 2019;102:728–731. doi: 10.1016/j.neuron.2019.03.026.normal* [[DOI](#)] [[PubMed](#)] [[Google Scholar](#)]
142. *Zambelli Z., Jakobsson C.E., Threadgokd L., Fidalgo A.R., Halstead E.J., Dimitriou D. Exploring the feasibility and acceptability of a sleep wearable headband among a community sample of chronic pain individuals: An at-home observational study. Digit. Health. 2022;8:20552076221097504. doi: 10.1177/20552076221097504.normal* [[DOI](#)] [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]
143. *De Fazio R., Mattei V., Al-Naami B., De Vittorio M. Methodologies and Wearable Devices to Monitor Biophysical Parameters Related to Sleep Dysfunctions: An Overview. Micromachines. 2022;13:1335. doi: 10.3390/mi13081335.normal* [[DOI](#)] [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]
144. *Cay G., Ravichandran V., Sadhu S., Zisk A.H. Recent advancement in sleep technologies: A literature review on clinical standards, sensors, apps, and AI methods. IEEE Access. 2022;99:104737–104756. doi: 10.1109/ACCESS.2022.3210518.normal* [[DOI](#)] [[Google](#)]

## [Scholar](#)]

145. Wei Z., Li W.Y., Wang X., Qin Z., Zhou J., Wang W. *Evaluation of a non-contact ultra-wideband bio-radar sleep monitoring device for screening of sleep breathing disease. Sleep Breath. 2022;26:689–696. doi: 10.1007/s11325-021-02424-x.* normal [[DOI](#)] [[PubMed](#)] [[Google Scholar](#)]
146. Zhang F., Zarkada G., Yi S., Eichmann A. *Lymphatic endothelial cell junctions: Molecular regulation in physiology and diseases. Front. Physiol. 2020;11:509. doi: 10.3389/fphys.2020.00509.* normal [[DOI](#)] [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]
147. Scallan J.P., Huxley V.H. *In vivo determination of collecting lymphatic vessel permeability to albumin: A role for lymphatics in exchange. J. Physiol. 2010;588:243–254. doi: 10.1113/jphysiol.2009.179622.* normal [[DOI](#)] [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]
148. Claesson-Welsh L., Dejana E., McDonald D.M. *Permeability of the endothelial barrier: Identifying and reconciling controversies. Trends Mol. Med. 2021;27:314–331. doi: 10.1016/j.molmed.2020.11.006.* normal [[DOI](#)] [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]
149. Kuan E.L., Ivanov S., Bridenbaugh E.A. *Collecting lymphatic vessel permeability facilitates adipose tissue inflammation and distribution of antigen to lymph node–homing adipose tissue dendritic cells. J. Immunol. 2015;194:5200–5210. doi: 10.4049/jimmunol.1500221.* normal [[DOI](#)] [[PMC free article](#)]

[\[PubMed\]](#) [\[Google Scholar\]](#)

150. Banerji S., Ni J., Wang S.-X., Clasper S., Su J., Tammi R., Jones M., Jackson D.G. Lyve-1, a new homologue of the cd44 glycoprotein, is a lymph-specific receptor for hyaluronan. *J. Cell Biol.* 1999;144:789–801. doi:

10.1083/jcb.144.4.789.normal [\[DOI\]](#) [\[PMC free article\]](#)

[\[PubMed\]](#) [\[Google Scholar\]](#)

151. Karu T.I., Pyatibrat L.V., Afanasyeva N.I. Cellular effects of low power laser therapy can be mediated by nitric oxide. *Lasers Surg. Med.* 2005;36:307–314. doi:

10.1002/lsm.20148.normal [\[DOI\]](#) [\[PubMed\]](#) [\[Google Scholar\]](#)

152. Murad F. Discovery of some of the biological effects of nitric oxide and its role in cell signaling. *Biosci. Rep.*

2004;24:452–474. doi: 10.1007/s10540-005-2741-8.normal

[\[DOI\]](#) [\[PubMed\]](#) [\[Google Scholar\]](#)

153. Colombo E., Signore A., Aicardi S., Zekiy A., Utyuzh A., Benedicenti S., Amaroli A. Experimental and clinical applications of red and near-infrared photobiomodulation on endothelial dysfunction: A review. *Biomedicines.* 2021;9:274. doi: 10.3390/biomedicines9030274.normal [\[DOI\]](#) [\[PMC free](#)

[article\]](#) [\[PubMed\]](#) [\[Google Scholar\]](#)

154. Uozumi Y., Nawashiro H., Sato S., Kawauchi S., Shima K., Kikuchi M. Targeted increase in cerebral blood flow by tran-scranial near-infrared laser irradiation. *Lasers Surg. Med.* 2010;42:566–576. doi: 10.1002/lsm.20938.normal

[\[DOI\]](#) [\[PubMed\]](#) [\[Google Scholar\]](#)

[\[DOI\]](#) [\[PubMed\]](#) [\[Google Scholar\]](#)

155. Litscher G., Min L., Passegger C.A., Litscher D., Li M.,

- Wang M., Ghaffari-Tabrizi-Wizsy N., Stelzer I., Feigl G., Gaischek I., et al. *Transcranial Yellow, Red, and Infrared Laser and LED Stimulation: Changes of Vascular Parameters in a Chick Embryo Model*. *Integr. Med. Int.* 2015;2:80–89. doi: 10.1159/000431176.normal [[DOI](#)] [[Google Scholar](#)]
156. Weihrauch D., Keszler A., Lindemer B., Krolikowski J., Lohr N.L. *Red light stimulates vasodilation through extracellular vesicle trafficking*. *J. Photochem. Photobiol. B Biol.* 2021;220:112212. doi: 10.1016/j.jphotobiol.2021.112212.normal [[DOI](#)] [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]
157. Scallan J.P., Zawieja S.D., Castorena-Gonzalez J., Davis M.J. *Lymphatic pumping: Mechanics, mechanisms and malfunction*. *J. Physiol.* 2016;594:5749–5768. doi: 10.1113/JP272088.normal [[DOI](#)] [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]
158. Stanley C.P., Maghzal G.J., Ayer A., Talib J., Giltrap A.M., Shengule S., Wolhuter K., Wang Y., Chadha P., Suarna C., et al. *Singlet molecular oxygen regulates vascular tone and blood pressure in inflammation*. *Nature*. 2019;566:548–552. doi: 10.1038/s41586-019-0947-3.normal [[DOI](#)] [[PubMed](#)] [[Google Scholar](#)]
159. Zakharov S.D., Ivanov A.V. *Light-oxygen effect in cells and its potential applications in tumour therapy (review)* *Quantum Electron.* 1999;29:1031–1053. doi: 10.1070/QE1999v029n12ABEH001629.normal [[DOI](#)] [[Google Scholar](#)]

160. Sokolovski S.G., Zolotovskaya S.A., Goltsov A., Pourreyron C., South A.P., Rafailov E.U. Infrared laser pulse triggers increased singlet oxygen production in tumour cells. *Sci. Rep.* 2013;3:3484. doi: 10.1038/srep03484.normal

[[DOI](#)] [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]

161. Khokhlova A., Zolotovskii I., Sokolovski S., Saenko Y., Rafailov E., Stoliarov D., Pogodina E., Svetukhin V., Sibirny V., Fotiadi A. The light-oxygen effect in biological cells enhanced by highly localized surface plasmon-polaritons.

*Sci. Rep.* 2019;9:18435. doi: 10.1038/s41598-019-54905-5.normal

[[DOI](#)] [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]

162. Khokhlova A., Zolotovskii I., Stoliarov D., Vorsina S., Liamina D., Pogodina E., Fotiadi A.A., Sokolovski S.G., Saenko Y., Rafailov E.U. The photobiomodulation of vital parameters of the cancer cell culture by low dose of near-IR laser irradiation. *IEEE J. Sel. Top. Quantum Electron.*

2019;25:1–10. doi: 10.1109/JSTQE.2018.2854539.normal

[[DOI](#)] [[Google Scholar](#)]

163. Ronsein G.E., Oliveira M.C., Miyamoto S., Medeiros M.H.G., Di Mascio P. Tryptophan oxidation by singlet molecular oxygen [O<sub>2</sub>(<sup>1</sup>Δ<sub>g</sub>): Mechanistic studies using <sup>18</sup>O-labeled hydroperoxides, mass spectrometry, and light emission measurements. *Chem. Res. Toxicol.* 2008;21:1271–

1283. doi: 10.1021/tx800026g.normal

[[DOI](#)] [[PubMed](#)] [[Google Scholar](#)]

164. Wang Y., Liu H., McKenzie G., Witting P.K., Stasch J.-P.,

Hahn M., Changsirivathanathamrong D., Wu B.J., Ball H.J., Thomas S.R., et al. Kynurenine is an endothelium-derived relaxing factor produced during inflammation. *Nat. Med.* 2010;16:279–285. doi: 10.1038/nm.2092.normal [[DOI](#)] [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]

165.Lane H. Cell biology: Power games. *Nature.* 2006;443:901–903. doi: 10.1038/443901a.normal [[DOI](#)] [[PubMed](#)] [[Google Scholar](#)]

166.Waypa G.B., Smith K.A., Schumacker P.T. O<sub>2</sub> sensing, mitochondria and ROS signaling: The fog is lifting. *Mol. Aspects Med.* 2016;47:76–89. doi: 10.1016/j.mam.2016.01.002.normal [[DOI](#)] [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]

167.Zhao Y., Vanhoutte P.M., Leung S.W. Vascular nitric oxide: Beyond eNOS. *J. Pharmacol. Sci.* 2015;129:83–94. doi: 10.1016/j.jphs.2015.09.002.normal [[DOI](#)] [[PubMed](#)] [[Google Scholar](#)]

168.Cassano P., Petrie S.P., Hamblin M.R., Henderson T.A., Iosifescu D.V. Review of transcranial photobiomodulation for major depressive disorder: Targeting brain metabolism, inflammation, oxidative stress, and neurogenesis. *Neurophot.* 2016;3:031404. doi: 10.1117/1.NPh.3.3.031404.normal [[DOI](#)] [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]

169.Zhang Z., Shen Q., Wu X., Zhang D., Xing D. Activation of PKA/SIRT1 signaling pathway by photobiomodulation therapy reduces: A levels in Alzheimer's disease models.

*Aging Cell.* 2020;19:e13054. doi: 10.1111/ace1.13054.normal

[[DOI](#)] [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]

170.Duggett N.A., Chazot P.L. Low-Intensity Light Therapy (1068 nm) Protects CAD Neuroblastoma Cells from

Amyloid-Mediated Cell Death. *Biol. Med.* 2014;1:2.normal

[[Google Scholar](#)]

171.Tao L., Liu Q., Zhang F., Fu Y., Zhu X., Weng X., Wei X.

Microglia modulation with 1070-nm light attenuates: A burden and cognitive impairment in Alzheimer's disease

mouse model. *Light Sci. Appl.* 2021;10:1–14. doi:

10.1038/s41377-021-00617-3.normal [[DOI](#)] [[PMC free](#)

[article](#)] [[PubMed](#)] [[Google Scholar](#)]

172.Yue X., Mei Y., Zhang Y., Tong Z., Cui D., Yang J., Wang

A., Wang R., Fei X., Ai L. New insight into Alzheimer's

disease: Light reverses: An obstructed interstitial fluid flow and ameliorates memory decline in APP/PS1 mice.

*Alzheimer's Dement. Transl. Res. Clin. Interv.* 2019;5:671–

684. doi: 10.1016/j.trci.2019.09.007.normal [[DOI](#)] [[PMC free](#)

[article](#)] [[PubMed](#)] [[Google Scholar](#)]

173.Lee G.-H., Kim H., Lee G.H., Kwon W., Too S., Myung D.,

Yun S.H., Bao Z., Hahn S.K. Multifunctional materials for

implantable and wearable photonic healthcare devices. *Nat.*

*Rev. Mater.* 2020;5:149–165. doi: 10.1038/s41578-019-

0167-3.normal [[DOI](#)] [[PMC free article](#)] [[PubMed](#)] [[Google](#)

[Scholar](#)]

174.Jinno H., Yokota Y., Koizumi M., Yukita W. Self-powered

ultraflexible photonic skin for continuous bio-signal

detection via air-operation-stable polymer light-emitting diodes. *Nat. Commun.* 2021;12:2234. doi: 10.1038/s41467-021-22558-6.normal [[DOI](#)] [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]

175.Da Mesquita S., Papadopoulos Z., Dykstra T., Brase L., Farias F., Wall M., Jiang H., Kodira C., de Lima K., Herz J., et al. Meningeal lymphatics affect microglia responses and anti-A $\beta$  immunotherapy. *Nature.* 2021;593:255–260. doi: 10.1038/s41586-021-03489-0.normal [[DOI](#)] [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]

176.Salloway S., Chalkias S., Barkhof F., Burkett P., Barakos J., Purcell D., Suhy J., Forrestal F., Tian Y., Umans K., et al. Amyloid-Related Imaging Abnormalities in 2 Phase 3 Studies Evaluating Aducanumab in Patients with Early Alzheimer Disease. *JAMA Neurol.* 2022;79:13–21. doi: 10.1001/jamaneurol.2021.4161.normal [[DOI](#)] [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]

177.Lustenberger C., Ferster M.L., Huwiler S., Brogli L., Werth E., Huber R., Karlen W. Auditory deep sleep stimulation in older adults at home: A randomized crossover trial. *Commun. Med.* 2022;2:30. doi: 10.1038/s43856-022-00096-6.normal [[DOI](#)] [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]

## Associated Data

*This section collects any data citations, data availability*

*statements, or supplementary materials included in this article.*

## **Data Availability Statement**

Not applicable.

Articles from International Journal of Molecular Sciences are provided here courtesy of **Multidisciplinary Digital Publishing Institute (MDPI)**